

Algorithms and Data Structures 2011/12
Week 10 tutorial solutions (Tues 22nd - Fri 25th Nov)

1. Show how to determine in $O(n^2 \lg n)$ time whether any three points in a set of n points are co-linear.

answer: This one depends on observing that *if* three points are co-linear, and we draw them in a line, then one of the three points must be lowest - ie, have the smallest y -value (if there are ties, we take the leftmost such point). Suppose p, q, r are co-linear, and p is the lowest. Then the polar angle of q wrt p is the same as the polar angle of r wrt p .

Our algorithm is as follows:

Choose every point p in turn, to act as the origin.

Then sort all of the other points in terms of their *polar angle* wrt p .

Finally do a linear search of the sorted points - if any two neighboring points in the sort have equal polar angles (mod π), they are co-linear with p , and we can terminate with **yes**.

Repeat until all points have been considered as the origin.

I did not yet explain how we can sort by polar angle - it is not too difficult. We can use a standard sorting algorithm such as MERGESORT, except that we must change the comparison operator. That can be done by observing that we can compare the polar angles of q, r wrt p by considering the vectors \vec{pq} and \vec{pr} . By definition, the vector \vec{pr} is anti-clockwise of \vec{pq} iff $(r - p) \times (q - p) < 0$. This is *almost* enough to order r and q by polar angle wrt p . First suppose the the y -coordinate of $q - p$ is non-negative:

If $(r - p) \times (q - p) < 0$, then r has greater polar angle wrt p than q ;

alternatively if $(r - p) \times (q - p) > 0$, then q has polar angle greater than r *if and only if* $r - p$ has a non-negative y -coordinate.

If $(r - p) \times (q - p) = 0$, then we need to check the y coordinate of $r - p$.

We can do similar case analysis for $q - p$ having a negative y -coordinate.

This polar-angle comparison operator can (clearly) be evaluated in constant time for any pair q, r . Hence by plugging the new comparison into MergeSort (say), we can sort polar angles wrt p in $\Theta(n \lg n)$ time. Finally in a linear walk through the array we can search for any pair of neighbouring points q, q' such that $(q - p) \times (q' - p) = 0$.

Notice that in the last phase (where we do a linear scan looking for neighbours with 0 cross product) we are forgetting to check for q, q' which are a polar angle π apart. This does not hurt our algorithm - even if it is the case that the 3 colinear points are such that p lies *between* q and q' , we will *observe* the co-linearity whenever we choose q or q' as our base point, whichever happens first.

2. Given a point $p_0 = (x_0, y_0)$, the *right horizontal ray* from p_0 is the set of points $\{p = (x, y_0) : x \geq x_0\}$, that is, it is the set of points due right of p_0 . Show how to determine whether a right horizontal ray from a given p_0 intersects a line segment $\overline{p_1 p_2}$ in $O(1)$ time, by reducing the problem to that of two line segments intersecting.

answer: This one is not too difficult. It depends on a simple observation - the line segment $\overline{p_1 p_2}$ is bounded in space. Hence if the right horizontal line from p_0 is to intersect $\overline{p_1 p_2}$, it must achieve this intersection within the ‘bounding box’ for $\overline{p_1 p_2}$. Let $p_0 = (x_0, y_0)$, $p_1 = (x_1, y_1)$, $p_2 = (x_2, y_2)$. We only need to consider the right horizontal line from p_0 between x_1 and x_2 . Hence we define $p_3 = (x_1, y_0)$ and $p_4 = (x_2, y_0)$. Then we can test for intersection of $\overline{p_1 p_2}$ with the horizontal ray by testing whether $\overline{p_1 p_2}$ intersects $\overline{p_3 p_4}$.

3. Show that there may be $\Theta(n^2)$ intersections in a set of n line segments.

answer: To see this, draw an n -by- n grid in the plane. Specifically we have:

- $n/2$ horizontal line segments of the form $((0, i), (n/2, i))$, for $1 \leq i \leq n/2$.
- $n/2$ vertical line segments of the form $((j, 0), (j, n/2))$, for $1 \leq j \leq n/2$.

It is clear from the definition that every horizontal line segment intersects every vertical line segment - moreover no more than two line segments intersect at any point. Hence there are $(n/2)^2 = \Theta(n^2)$ intersections in total.

4. In the *online convex hull problem*, we are given the set Q of n points one point at a time. After receiving each point, we are to compute the convex hull of the points seen so far. Obviously, we could run Graham’s scan once for each point, with a total running time of $O(n^2 \lg n)$. Show how to improve this slightly, by showing we can solve the online convex hull problem in $O(n^2)$.

This is Ex. 33.3-5 of [CLRS]. Ex. 35.3-5 of [CLR].

answer: It is good to first think about why the naïve algorithm (re-running Graham’s scan each time a point is added) will only lead to the bound $O(n^2 \lg n)$. This is because, if Graham’s scan is $O(n \lg n)$, this means there is some $c \geq 0$ such that the running time of Graham’s scan is $\leq cn \lg n$ for sufficiently large n . This gives us the following upper bound

$$\sum_{k=3}^n ck \lg n$$

of the running time of the naïve online algorithm. Taking only the terms from $k = n/2$ to $k = n$, this would give us an expression as large as $\frac{n^2}{4}(\lg n - 1)$ which is of the form $\Theta(n^2 \lg n)$. So we need to work harder...

In coming up with our better online algorithm, note that the points p_1, \dots, p_n may arrive in any order, the ordering of them does not imply any geometric relationship

(it is important to remember this, to distinguish from the scenario of the standard Convex Hull algorithm, where we sort all points together first, and then access points in order of polar angle)

For our improved algorithm, we assume, that the convex hull of a set of points is *represented* by its points presented in *anti-clockwise* order, starting with the *bottom-most* point (if there is more than one point with this y -coordinate, then take the left-most of these). In the online setting, we will consider a number of such “convex hulls”, as points are added.


The naïve algorithm was defined by running the convex hull algorithm again every time a new point was added - this having the time-bound $O(k \lg(k))$ for the update for the k th point.

Our better algorithm will just use $O(k)$ work to update the convex hull for the k -th point. We will not *need* to do the sorting of points (which is what drives the $O(n \lg(n))$ running time of Graham’s scan) when we are just adding a new extra point to a current convex hull. This is because if we have the convex hull $C(k-1)$ of the points $\{p_1, \dots, p_{k-1}\}$, then this lists the points of that hull in counterclockwise order from the bottom-most (and left-most if breaking ties) point in that set.

Now consider the work we need to do in updating the convex hull $C(k-1)$ to obtain $C(k)$, ie to consider the change on adding p_k .

- First suppose p_k sits in the interior of $C(k-1)$. Then $C(k)$ is equal to $C(k-1)$. We can detect this case by the fact that if $p_{1,k-1}, \dots, p_{\ell(k-1),k-1}$ are the points of $C(k-1)$ in anti-clockwise order from the lowest one $p_{1,k-1}$, then for every $1 \leq i \leq \ell(k-1)$, $p_{i,k-1} \rightarrow p_{i+1,k-1} \rightarrow p_k$ involves a *left-turn* (note we perform $i+1$ in a wrap around fashion, with $p_{\ell(k-1)+1,k-1} =_{\text{def}} p_{1,k-1}$)
- Alternatively suppose that p_k is on the exterior of $C(k-1)$. Then the convex hull consists of p_k plus one continuous segment from $C(k-1)$ - usually this segment will consist of most of the existing points on the convex hull $C(k-1)$. In this scenario, the left/right turns of the convex hull $C(k-1)$ wrt p_k can be characterised as follows:
 - In considering the points of $C(k-1)$ from the bottom-most point $p_{1,k-1}$ onwards, let i be the first point such that $p_{i,k-1} \rightarrow p_{i+1,k-1} \rightarrow p_k$ involves a *right turn*.
 - Subsequent to finding the i defined above, let j be the first index *after* i such that $p_{j,k-1} \rightarrow p_{j+1,k-1} \rightarrow p_k$ involves a *left turn*.
 - It is easy to argue that $p_{h,k-1} \rightarrow p_{h+1,k-1} \rightarrow p_k$ is always a left turn from $h = j$ all the way round to $h = i-1$.
 - The convex hull $C(k)$ is equal to $p_{1,k-1} \dots, p_{i,k-1}, p_k, p_j, p_{j+1,k-1} \dots, p_{\ell(k-1),k-1}$.

The following picture is a good reference point for the discussion.



Now we observe that the updating of $C(k-1)$ to become $C(k)$ just involves a linear scan of the existing points (of which there are at most $k-1$) on $C(k-1)$, with the “work done” in considering each point, being just constant-time (doing the left-turn/right-turn test). Overall the update for the new point p_k takes $O(n)$ time.

5. Prove that the problem of finding the Convex Hull of n points has a lower bound of $\Omega(n \lg n)$. For this, think about using a *reduction* from sorting to Convex Hull (that is, think about how to use a Convex Hull algorithm to sort a list of numbers).

answer: The best approach for this one is to think about *what* the convex hull problem is.

The question is asking for a *reduction* - in other words, showing how to map one problem to another.

There is only one observation needed - if we are given a set of points which form a convex polytope in the plane, then the convex hull is just the list of those points in counterclockwise order (guess this last point is important, the convex hull lists the points in order).

We are given a list of numbers which we wish to sort. I’m not going to assume anything special about them, no bounds on size, they can be reals (though not complex), etc. However as we know from our information-theoretic lower bounds on sorting, we may assume that the numbers we wish to sort are *distinct* (this is a special case of sorting which we know has complexity $\Omega(n \lg n)$)

For our reduction, we first do a linear scan of the numbers, separating them into positive and negative sets. We will do two rounds of the convex hull problem.

First take the set of non-negative numbers:

We will map each number to a point in the plane. To do this I want to think of a mapping which will arrange the points along a convex curve in the plane (so that all points will wind up on the convex hull). So I will work with the curve $x \rightarrow x^2$. For every non-negative number x in our list of elements to be sorted, we add (x, x^2) to the set of points for convex hull. Finally, we add an extra point $(-1, 0)$ (to act as an anchor). Note that because the curve $x \rightarrow x^2$ is convex, *every* point (x, x^2) will end up on the convex hull. Also $(-1, 0)$ will lie on the convex hull. Moreover, remember our requirement that the convex hull points should be output in anti-clockwise order.

Hence the points will be returned in increasing order of (x, x^2) . Since this order is the same as the increasing order of the numbers x , we have sorted all positive numbers.

Similarly, we take the set of negative numbers and map every x here to $(|x|, x^2)$. Then after making a call to convex hull, the points are returned in increasing order of $|x|$, ie in decreasing order of x . We invert the list of sorted points, and take $-|x|$ for each point. Then we have the negative numbers in sorted order.

Finally we merge the two lists in $\Theta(n)$ time.

Hence we have an algorithm which uses two calls to the Convex Hull problem and a linear amount of “extra work” (eg, partitioning into small/large, mapping from numbers to points and back again, merging the two sorted lists) to sort the list of input numbers. We can write

$$T_{\text{sort}}(n) \leq 2T_{\text{hull}}(n) + \Theta(n).$$

However we know sorting is $\Omega(n \lg n)$. Hence $2T_{\text{hull}}(n) + \Theta(n)$ is $\Omega(n \lg n)$. We know $\Theta(n)$ cannot be $\Omega(n \lg n)$, hence it must be the case that $2T_{\text{hull}}(n)$ is $\Omega(n \lg n)$.

note: This last step is why we need to count up the “extra work” done on top of the calls to the convex-hull problem (we have to arrange things so that the lower bound must be satisfied by the convex hull calls).

note 2: We cheated a *tiny* bit, but I think this is a nice question and worth doing anyhow. If we are being strict with ourselves we should not assume $\Omega(n \lg n)$ for this version of sorting. We proved $\Omega(n \lg n)$ for *comparison-based sorting*, and here we have a bit more information, we know the elements to be sorted are at least real numbers.

Strictly we can't be sure we have an $\Omega(n \lg n)$ bound for this case - though on the other hand, Radix sort does not apply to reals in general (only reals which can be represented with a bounded number of bits). The $\Omega(n \lg(n))$ lower bound is a sensible thing to believe for inputs which are general reals, though we don't have an answer one way or the other.

Mary Cryan