

Discrete Mathematics & Mathematical Reasoning

Basic Structures: Sets, Functions, Relations, Sequences and Sums

Colin Stirling

Informatics

Sets

- A set is an unordered collection of elements

Sets

- A set is an unordered collection of elements
- $A = \{3, 2, 1, 0\} = \{1, 2, 0, 3\}$

Sets

- A set is an unordered collection of elements
- $A = \{3, 2, 1, 0\} = \{1, 2, 0, 3\}$
- Membership $3 \in A$
- Non-membership $5 \notin A$

Sets

- A set is an unordered collection of elements
- $A = \{3, 2, 1, 0\} = \{1, 2, 0, 3\}$
- Membership $3 \in A$
- Non-membership $5 \notin A$
- Emptyset $\emptyset = \{ \}$

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ Natural numbers

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ **Natural numbers**

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integers

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ Natural numbers

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integers

$\mathbb{Z}^+ = \{z \in \mathbb{Z} \mid z > 0\}$ Positive integers

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ Natural numbers

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integers

$\mathbb{Z}^+ = \{z \in \mathbb{Z} \mid z > 0\}$ Positive integers

\mathbb{R} Real numbers

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ Natural numbers

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integers

$\mathbb{Z}^+ = \{z \in \mathbb{Z} \mid z > 0\}$ Positive integers

\mathbb{R} Real numbers

$\mathbb{R}^+ = \{r \in \mathbb{R} \mid r > 0\}$ Positive real numbers

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ Natural numbers

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integers

$\mathbb{Z}^+ = \{z \in \mathbb{Z} \mid z > 0\}$ Positive integers

\mathbb{R} Real numbers

$\mathbb{R}^+ = \{r \in \mathbb{R} \mid r > 0\}$ Positive real numbers

$\mathbb{Q} = \{\frac{a}{b} \mid a \in \mathbb{Z}, b \in \mathbb{Z}^+\}$ Rational numbers

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ Natural numbers

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integers

$\mathbb{Z}^+ = \{z \in \mathbb{Z} \mid z > 0\}$ Positive integers

\mathbb{R} Real numbers

$\mathbb{R}^+ = \{r \in \mathbb{R} \mid r > 0\}$ Positive real numbers

$\mathbb{Q} = \{\frac{a}{b} \mid a \in \mathbb{Z}, b \in \mathbb{Z}^+\}$ Rational numbers

$\mathbb{Q}^+ = \{\frac{a}{b} \mid a \in \mathbb{Z}^+, b \in \mathbb{Z}^+\}$ Positive rational numbers

Some important sets (boldface in the textbook)

$\mathbb{B} = \{\text{true}, \text{false}\}$ Boolean values

$\mathbb{N} = \{0, 1, 2, 3, \dots\}$ Natural numbers

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integers

$\mathbb{Z}^+ = \{z \in \mathbb{Z} \mid z > 0\}$ Positive integers

\mathbb{R} Real numbers

$\mathbb{R}^+ = \{r \in \mathbb{R} \mid r > 0\}$ Positive real numbers

$\mathbb{Q} = \{\frac{a}{b} \mid a \in \mathbb{Z}, b \in \mathbb{Z}^+\}$ Rational numbers

$\mathbb{Q}^+ = \{\frac{a}{b} \mid a \in \mathbb{Z}^+, b \in \mathbb{Z}^+\}$ Positive rational numbers

\mathbb{C} Complex numbers

Sets defined using comprehension

- $S = \{x \mid P(x)\}$ where $P(x)$ is a predicate

Sets defined using comprehension

- $S = \{x \mid P(x)\}$ where $P(x)$ is a predicate
- $\mathbb{Z}^+, \mathbb{R}^+, \mathbb{Q}^+$

Sets defined using comprehension

- $S = \{x \mid P(x)\}$ where $P(x)$ is a predicate
- $\mathbb{Z}^+, \mathbb{R}^+, \mathbb{Q}^+$
- $\{x \mid x \in \mathbb{N} \wedge 2 \text{ divides } x\} = \{x \in \mathbb{N} \mid \exists k(x = 2k)\}$

Sets defined using comprehension

- $S = \{x \mid P(x)\}$ where $P(x)$ is a predicate
- $\mathbb{Z}^+, \mathbb{R}^+, \mathbb{Q}^+$
- $\{x \mid x \in \mathbb{N} \wedge 2 \text{ divides } x\} = \{x \in \mathbb{N} \mid \exists k(x = 2k)\}$
- Closed intervals $[0, 1] = \{r \mid 0 \leq r \leq 1\}$

Notation

- $A \cup B$ union; $A \cap B$ intersection

Notation

- $A \cup B$ union; $A \cap B$ intersection
- $A - B$ difference; \bar{A} complement

Notation

- $A \cup B$ union; $A \cap B$ intersection
- $A - B$ difference; \bar{A} complement
- If A_i are sets for all $i \in I$ then $\bigcup_{i \in I} A_i$ and $\bigcap_{i \in I} A_i$ are sets

Notation

- $A \cup B$ union; $A \cap B$ intersection
- $A - B$ difference; \bar{A} complement
- If A_i are sets for all $i \in I$ then $\bigcup_{i \in I} A_i$ and $\bigcap_{i \in I} A_i$ are sets
- $A \subseteq B$ subset; $A \supseteq B$ superset

Notation

- $A \cup B$ union; $A \cap B$ intersection
- $A - B$ difference; \bar{A} complement
- If A_i are sets for all $i \in I$ then $\bigcup_{i \in I} A_i$ and $\bigcap_{i \in I} A_i$ are sets
- $A \subseteq B$ subset; $A \supseteq B$ superset
- $A = B$ set equality

Notation

- $A \cup B$ union; $A \cap B$ intersection
- $A - B$ difference; \bar{A} complement
- If A_i are sets for all $i \in I$ then $\bigcup_{i \in I} A_i$ and $\bigcap_{i \in I} A_i$ are sets
- $A \subseteq B$ subset; $A \supseteq B$ superset
- $A = B$ set equality
- $\mathcal{P}(A)$ powerset (set of all subsets of A); also 2^A

Notation

- $A \cup B$ union; $A \cap B$ intersection
- $A - B$ difference; \bar{A} complement
- If A_i are sets for all $i \in I$ then $\bigcup_{i \in I} A_i$ and $\bigcap_{i \in I} A_i$ are sets
- $A \subseteq B$ subset; $A \supseteq B$ superset
- $A = B$ set equality
- $\mathcal{P}(A)$ powerset (set of all subsets of A); also 2^A
- $|A|$ cardinality

Notation

- $A \cup B$ union; $A \cap B$ intersection
- $A - B$ difference; \bar{A} complement
- If A_i are sets for all $i \in I$ then $\bigcup_{i \in I} A_i$ and $\bigcap_{i \in I} A_i$ are sets
- $A \subseteq B$ subset; $A \supseteq B$ superset
- $A = B$ set equality
- $\mathcal{P}(A)$ powerset (set of all subsets of A); also 2^A
- $|A|$ cardinality
- $A \times B$ cartesian product (tuple sets)

Common set identities

- $A \cup A = A$ $A \cap A = A$

Common set identities

- $A \cup A = A$ $A \cap A = A$
- $\overline{\overline{A}} = A$

Common set identities

- $A \cup A = A$ $A \cap A = A$
- $\overline{\overline{A}} = A$
- $A \cup B = B \cup A$ $A \cap B = B \cap A$

Common set identities

- $A \cup A = A$ $A \cap A = A$
- $\overline{\overline{A}} = A$
- $A \cup B = B \cup A$ $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$

Common set identities

- $A \cup A = A$ $A \cap A = A$
- $\overline{\overline{A}} = A$
- $A \cup B = B \cup A$ $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $\overline{A \cap B} = \overline{A} \cup \overline{B}$ $\overline{A \cup B} = \overline{A} \cap \overline{B}$

Common set identities

- $A \cup A = A$ $A \cap A = A$
- $\overline{\overline{A}} = A$
- $A \cup B = B \cup A$ $A \cap B = B \cap A$
- $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ $A \cap (B \cup C) = (A \cap B) \cup (A \cap C)$
- $\overline{A \cap B} = \overline{A} \cup \overline{B}$ $\overline{A \cup B} = \overline{A} \cap \overline{B}$
- $A \cup (A \cap B) = A$ $A \cap (A \cup B) = A$

A proper mathematical definition of set is complicated (Russell's paradox)

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)
- The set of non-cats (all things that are not cats) is a member of itself

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)
- The set of non-cats (all things that are not cats) is a member of itself
- Let S be the set of all sets which are not members of themselves

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)
- The set of non-cats (all things that are not cats) is a member of itself
- Let S be the set of all sets which are not members of themselves
- $S = \{x \mid x \notin x\}$ (using naive comprehension)

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)
- The set of non-cats (all things that are not cats) is a member of itself
- Let S be the set of all sets which are not members of themselves
- $S = \{x \mid x \notin x\}$ (using naive comprehension)
- Question: is S a member of itself ($S \in S$) ?

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)
- The set of non-cats (all things that are not cats) is a member of itself
- Let S be the set of all sets which are not members of themselves
- $S = \{x \mid x \notin x\}$ (using naive comprehension)
- Question: is S a member of itself ($S \in S$) ?
- $S \in S$ provided that $S \notin S$; $S \notin S$ provided that $S \in S$

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)
- The set of non-cats (all things that are not cats) is a member of itself
- Let S be the set of all sets which are not members of themselves
- $S = \{x \mid x \notin x\}$ (using naive comprehension)
- Question: is S a member of itself ($S \in S$) ?
- $S \in S$ provided that $S \notin S$; $S \notin S$ provided that $S \in S$
- There cannot be such a set S

A proper mathematical definition of set is complicated (Russell's paradox)

- The set of cats is not a cat (is not a member of itself)
- The set of non-cats (all things that are not cats) is a member of itself
- Let S be the set of all sets which are not members of themselves
- $S = \{x \mid x \notin x\}$ (using naive comprehension)
- Question: is S a member of itself ($S \in S$) ?
- $S \in S$ provided that $S \notin S$; $S \notin S$ provided that $S \in S$
- There cannot be such a set S
- Modern formulations (such as Zermelo-Fraenkel set theory) restrict comprehension. (However, it is impossible to prove in ZF that ZF is consistent unless ZF is inconsistent.)

Functions

- Assume A and B are non-empty sets

Functions

- Assume A and B are non-empty sets
- f is a function from A to B if f assigns to each element of A a unique element of B

Functions

- Assume A and B are non-empty sets
- f is a function from A to B if f assigns to each element of A a unique element of B
- Write $f(a) = b$ if f assigns b to a

Functions

- Assume A and B are non-empty sets
- f is a function from A to B if f assigns to each element of A a unique element of B
- Write $f(a) = b$ if f assigns b to a
- $f : A \rightarrow B$ if f is a function from A to B

Functions

- Assume A and B are non-empty sets
- f is a function from A to B if f assigns to each element of A a unique element of B
- Write $f(a) = b$ if f assigns b to a
- $f : A \rightarrow B$ if f is a function from A to B
- If $f : A \rightarrow B$, A is the domain and B is codomain (range)

Examples

- $f : \text{DMMR_Students} \rightarrow \text{Percentages}$

Examples

- $f : \text{DMMR_Students} \rightarrow \text{Percentages}$
- $\iota_A : A \rightarrow A$ where $\iota_A(a) = a$ identity

Examples

- $f : \text{DMMR_Students} \rightarrow \text{Percentages}$
- $\iota_A : A \rightarrow A$ where $\iota_A(a) = a$ identity
- $\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$: floor largest integer less than or equal to x

Examples

- $f : \text{DMMR_Students} \rightarrow \text{Percentages}$
- $\iota_A : A \rightarrow A$ where $\iota_A(a) = a$ identity
- $\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$: floor largest integer less than or equal to x
what are $\lfloor \frac{1}{2} \rfloor$ $\lfloor -6.1 \rfloor$

Examples

- $f : \text{DMMR_Students} \rightarrow \text{Percentages}$
- $\iota_A : A \rightarrow A$ where $\iota_A(a) = a$ identity
- $\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$: floor largest integer less than or equal to x
what are $\lfloor \frac{1}{2} \rfloor$ $\lfloor -6.1 \rfloor$
- $\lceil x \rceil : \mathbb{R} \rightarrow \mathbb{Z}$: ceiling smallest integer greater than or equal to x

Examples

- $f : \text{DMMR_Students} \rightarrow \text{Percentages}$
- $\iota_A : A \rightarrow A$ where $\iota_A(a) = a$ identity
- $\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$: floor largest integer less than or equal to x
what are $\lfloor \frac{1}{2} \rfloor$ $\lfloor -6.1 \rfloor$
- $\lceil x \rceil : \mathbb{R} \rightarrow \mathbb{Z}$: ceiling smallest integer greater than or equal to x
what are $\lceil \frac{1}{2} \rceil$ $\lceil -6.1 \rceil$

Examples

- $f : \text{DMMR_Students} \rightarrow \text{Percentages}$
- $\iota_A : A \rightarrow A$ where $\iota_A(a) = a$ identity
- $\lfloor x \rfloor : \mathbb{R} \rightarrow \mathbb{Z}$: floor largest integer less than or equal to x
what are $\lfloor \frac{1}{2} \rfloor$ $\lfloor -6.1 \rfloor$
- $\lceil x \rceil : \mathbb{R} \rightarrow \mathbb{Z}$: ceiling smallest integer greater than or equal to x
what are $\lceil \frac{1}{2} \rceil$ $\lceil -6.1 \rceil$
- $! : \mathbb{N} \rightarrow \mathbb{N}$ Factorial
 $0! = 1$ $n! = 1 \cdot 2 \cdot \dots \cdot (n-1) \cdot n$ for $n > 0$

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective?

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective?

YES

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective?
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective?

YES

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective? YES

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective? YES
- Is the squaring function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ injective?

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective? YES
- Is the squaring function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ injective? NO

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective? YES
- Is the squaring function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ injective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ injective?

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective? YES
- Is the squaring function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ injective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ injective? NO

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective? YES
- Is the squaring function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ injective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ injective? NO
- Assume $m > 1$. Is $\text{mod } m : \mathbb{Z} \rightarrow \{0, \dots, m - 1\}$ injective?

One-to-one or injective functions

Definition

$f : A \rightarrow B$ is injective iff $\forall a, c \in A$ (if $f(a) = f(c)$ then $a = c$)

- Is the identity function $\iota_A : A \rightarrow A$ injective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ injective? YES
- Is the squaring function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ injective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ injective? NO
- Assume $m > 1$. Is $\text{mod } m : \mathbb{Z} \rightarrow \{0, \dots, m-1\}$ injective? NO

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective?

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective?

YES

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective?
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective?

YES

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective? NO

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective? NO
- Is the function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ surjective?

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective? NO
- Is the function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ surjective? NO

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective? NO
- Is the function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ surjective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ surjective?

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective? NO
- Is the function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ surjective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ surjective? NO

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective? NO
- Is the function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ surjective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ surjective? NO
- Assume $m > 1$. Is $\text{mod } m : \mathbb{Z} \rightarrow \{0, \dots, m - 1\}$ surjective?

Onto or surjective functions

Definition

$f : A \rightarrow B$ is surjective iff $\forall b \in B \exists a \in A (f(a) = b)$

- Is the identity function $\iota_A : A \rightarrow A$ surjective? YES
- Is the function $\sqrt{\cdot} : \mathbb{Z}^+ \rightarrow \mathbb{R}^+$ surjective? NO
- Is the function $\cdot^2 : \mathbb{Z} \rightarrow \mathbb{Z}$ surjective? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ surjective? NO
- Assume $m > 1$. Is $\text{mod } m : \mathbb{Z} \rightarrow \{0, \dots, m - 1\}$ surjective? YES

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection?

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection?

YES

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection?
- Is the function $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a bijection?

YES

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection? YES
- Is the function $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a bijection? YES

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection? YES
- Is the function $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a bijection? YES
- Is the function $\cdot^2 : \mathbb{R} \rightarrow \mathbb{R}$ a bijection?

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection? YES
- Is the function $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a bijection? YES
- Is the function $\cdot^2 : \mathbb{R} \rightarrow \mathbb{R}$ a bijection? NO

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection? YES
- Is the function $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a bijection? YES
- Is the function $\cdot^2 : \mathbb{R} \rightarrow \mathbb{R}$ a bijection? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ a bijection?

One-to-one correspondence or bijection

Definition

$f : A \rightarrow B$ is a bijection iff it is both injective and surjective

- Is the identity function $\iota_A : A \rightarrow A$ a bijection? YES
- Is the function $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ a bijection? YES
- Is the function $\cdot^2 : \mathbb{R} \rightarrow \mathbb{R}$ a bijection? NO
- Is the function $|\cdot| : \mathbb{R} \rightarrow \mathbb{R}$ a bijection? NO

Function composition

Definition

Let $f : B \rightarrow C$ and $g : A \rightarrow B$. The composition function $f \circ g : A \rightarrow C$ is
 $(f \circ g)(a) = f(g(a))$

Results about function composition

Theorem

The composition of two functions is a function

Results about function composition

Theorem

The composition of two functions is a function

Theorem

The composition of two injective functions is an injective function

Results about function composition

Theorem

The composition of two functions is a function

Theorem

The composition of two injective functions is an injective function

Theorem

The composition of two surjective functions is a surjective function

Results about function composition

Theorem

The composition of two functions is a function

Theorem

The composition of two injective functions is an injective function

Theorem

The composition of two surjective functions is a surjective function

Corollary

The composition of two bijections is a bijection

Inverse function


Definition

If $f : A \rightarrow B$ is a bijection, then the inverse of f , written $f^{-1} : B \rightarrow A$ is $f^{-1}(b) = a$ iff $f(a) = b$

Inverse function

Definition


If $f : A \rightarrow B$ is a bijection, then the inverse of f , written $f^{-1} : B \rightarrow A$ is $f^{-1}(b) = a$ iff $f(a) = b$



Inverse function

Definition

If $f : A \rightarrow B$ is a bijection, then the inverse of f , written $f^{-1} : B \rightarrow A$ is $f^{-1}(b) = a$ iff $f(a) = b$




What is the inverse of $\iota_A : A \rightarrow A$?

Inverse function

Definition

If $f : A \rightarrow B$ is a bijection, then the inverse of f , written $f^{-1} : B \rightarrow A$ is $f^{-1}(b) = a$ iff $f(a) = b$




What is the inverse of $\iota_A : A \rightarrow A$?

What is the inverse of $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$?

Inverse function

Definition

If $f : A \rightarrow B$ is a bijection, then the inverse of f , written $f^{-1} : B \rightarrow A$ is $f^{-1}(b) = a$ iff $f(a) = b$



What is the inverse of $\iota_A : A \rightarrow A$?

What is the inverse of $\sqrt{\cdot} : \mathbb{R}^+ \rightarrow \mathbb{R}^+$?

What is $f^{-1} \circ f$? and $f \circ f^{-1}$?

Relations

Definition

A binary relation R on sets A and B is a subset $R \subseteq A \times B$

Relations

Definition

A binary relation R on sets A and B is a subset $R \subseteq A \times B$

- R is a set of tuples (a, b) with $a \in A$ and $b \in B$

Relations

Definition

A binary relation R on sets A and B is a subset $R \subseteq A \times B$

- R is a set of tuples (a, b) with $a \in A$ and $b \in B$
- Often we write $a R b$ for $(a, b) \in R$

Relations

Definition

A binary relation R on sets A and B is a subset $R \subseteq A \times B$

- R is a set of tuples (a, b) with $a \in A$ and $b \in B$
- Often we write $a R b$ for $(a, b) \in R$
- A function f is a restricted relation where

$$\forall a \in A \exists b \in B (((a, b) \in f) \wedge \forall c \in B ((a, c) \in f \rightarrow c = b))$$

Relations

Definition

A binary relation R on sets A and B is a subset $R \subseteq A \times B$

- R is a set of tuples (a, b) with $a \in A$ and $b \in B$
- Often we write $a R b$ for $(a, b) \in R$
- A function f is a restricted relation where

$$\forall a \in A \exists b \in B (((a, b) \in f) \wedge \forall c \in B ((a, c) \in f \rightarrow c = b))$$

- R is a relation on A if $B = A$ (so, $R \subseteq A \times A$)

Relations

Definition

A binary relation R on sets A and B is a subset $R \subseteq A \times B$

- R is a set of tuples (a, b) with $a \in A$ and $b \in B$
- Often we write $a R b$ for $(a, b) \in R$
- A function f is a restricted relation where

$$\forall a \in A \exists b \in B (((a, b) \in f) \wedge \forall c \in B ((a, c) \in f \rightarrow c = b))$$

- R is a relation on A if $B = A$ (so, $R \subseteq A \times A$)

Relations

Definition

A binary relation R on sets A and B is a subset $R \subseteq A \times B$

- R is a set of tuples (a, b) with $a \in A$ and $b \in B$
- Often we write $a R b$ for $(a, b) \in R$
- A function f is a restricted relation where

$$\forall a \in A \exists b \in B (((a, b) \in f) \wedge \forall c \in B ((a, c) \in f \rightarrow c = b))$$

- R is a relation on A if $B = A$ (so, $R \subseteq A \times A$)

Definition

Given sets A_1, \dots, A_n a subset $R \subseteq A_1 \times \dots \times A_n$ is an n -ary relation

Examples

- $R \subseteq A \times B$, A students, B courses; $(\text{Colin}, \text{DMMR}) \in R$

Examples

- $R \subseteq A \times B$, A students, B courses; $(\text{Colin}, \text{DMMR}) \in R$
- **Graphs are relations on vertices: covered later in course**

Examples

- $R \subseteq A \times B$, A students, B courses; $(\text{Colin}, \text{DMMR}) \in R$
- **Graphs are relations on vertices: covered later in course**
- Divides $|: \mathbb{Z} \times \mathbb{Z}$ is $\{(a, b) \mid a \neq 0 \text{ and } \exists c \in \mathbb{Z} (b = ac)\}$

Examples

- $R \subseteq A \times B$, A students, B courses; (Colin, DMMR) $\in R$
- **Graphs are relations on vertices: covered later in course**
- Divides $| : \mathbb{Z} \times \mathbb{Z}$ is $\{(a, b) \mid a \neq 0 \text{ and } \exists c \in \mathbb{Z} (b = ac)\}$
- $R = \{(a, b) \mid m \text{ divides } a - b\}$ where $m > 1$ is an integer

Examples

- $R \subseteq A \times B$, A students, B courses; $(\text{Colin}, \text{DMMR}) \in R$
- **Graphs are relations on vertices: covered later in course**
- Divides $| : \mathbb{Z} \times \mathbb{Z}$ is $\{(a, b) \mid a \neq 0 \text{ and } \exists c \in \mathbb{Z} (b = ac)\}$
- **$R = \{(a, b) \mid m \text{ divides } a - b\}$ where $m > 1$ is an integer** Written as $a \equiv b \pmod{m}$

Notation

- $R \cup S$ union; $R \cap S$ intersection;

Notation

- $R \cup S$ union; $R \cap S$ intersection;
- If R_i are relations on $A \times B$ for all $i \in I$ then $\bigcup_{i \in I} R_i$ and $\bigcap_{i \in I} R_i$ are relations on $A \times B$

Notation

- $R \cup S$ union; $R \cap S$ intersection;
- If R_i are relations on $A \times B$ for all $i \in I$ then $\bigcup_{i \in I} R_i$ and $\bigcap_{i \in I} R_i$ are relations on $A \times B$
- $R \subseteq S$ subset and $R = S$ equality

Relation composition

Definition

Let $R \subseteq B \times C$ and $S \subseteq A \times B$. The composition relation $(R \circ S) \subseteq A \times C$ is $\{(a, c) \mid \exists b (a, b) \in S \wedge (b, c) \in R\}$

Relation composition

Definition

Let $R \subseteq B \times C$ and $S \subseteq A \times B$. The composition relation $(R \circ S) \subseteq A \times C$ is $\{(a, c) \mid \exists b (a, b) \in S \wedge (b, c) \in R\}$

Closure R is a relation on A :

- R^0 is the identity relation (ι_A)
- $R^{n+1} = R^n \circ R$
- $R^* = \bigcup_{n \geq 0} R^n$

Relation composition

Definition

Let $R \subseteq B \times C$ and $S \subseteq A \times B$. The composition relation $(R \circ S) \subseteq A \times C$ is $\{(a, c) \mid \exists b (a, b) \in S \wedge (b, c) \in R\}$

Closure R is a relation on A :

- R^0 is the identity relation (ι_A)
- $R^{n+1} = R^n \circ R$
- $R^* = \bigcup_{n \geq 0} R^n$

Example: reachability in a graph

Properties of binary relation R on A

- reflexive iff $\forall x \in A (x, x) \in R$

Properties of binary relation R on A

- reflexive iff $\forall x \in A (x, x) \in R$
- \leq , $=$, and $|$ are reflexive, but $<$ is not

Properties of binary relation R on A

- reflexive iff $\forall x \in A (x, x) \in R$
- \leq , $=$, and $|$ are reflexive, but $<$ is not
- symmetric iff $\forall x, y \in A ((x, y) \in R \rightarrow (y, x) \in R)$
- $=$ is symmetric, but \leq , $<$, and $|$ are not

Properties of binary relation R on A

- reflexive iff $\forall x \in A (x, x) \in R$
- \leq , $=$, and $|$ are reflexive, but $<$ is not
- symmetric iff $\forall x, y \in A ((x, y) \in R \rightarrow (y, x) \in R)$
- $=$ is symmetric, but \leq , $<$, and $|$ are not
- antisymmetric iff $\forall x, y \in A (((x, y) \in R \wedge (y, x) \in R) \rightarrow x = y)$

Properties of binary relation R on A

- reflexive iff $\forall x \in A (x, x) \in R$
- \leq , $=$, and $|$ are reflexive, but $<$ is not
- symmetric iff $\forall x, y \in A ((x, y) \in R \rightarrow (y, x) \in R)$
- $=$ is symmetric, but \leq , $<$, and $|$ are not
- antisymmetric iff $\forall x, y \in A (((x, y) \in R \wedge (y, x) \in R) \rightarrow x = y)$
- \leq , $=$, $<$ are antisymmetric but $|$ is not

Properties of binary relation R on A

- reflexive iff $\forall x \in A (x, x) \in R$
- \leq , $=$, and $|$ are reflexive, but $<$ is not
- symmetric iff $\forall x, y \in A ((x, y) \in R \rightarrow (y, x) \in R)$
- $=$ is symmetric, but \leq , $<$, and $|$ are not
- antisymmetric iff $\forall x, y \in A (((x, y) \in R \wedge (y, x) \in R) \rightarrow x = y)$
- \leq , $=$, $<$ are antisymmetric but $|$ is not
- transitive iff $\forall x, y, z \in A (((x, y) \in R \wedge (y, z) \in R) \rightarrow (x, z) \in R)$
- \leq , $=$, $<$, and $|$ are transitive

Properties of binary relation R on A

- reflexive iff $\forall x \in A (x, x) \in R$
- \leq , $=$, and $|$ are reflexive, but $<$ is not
- symmetric iff $\forall x, y \in A ((x, y) \in R \rightarrow (y, x) \in R)$
- $=$ is symmetric, but \leq , $<$, and $|$ are not
- antisymmetric iff $\forall x, y \in A (((x, y) \in R \wedge (y, x) \in R) \rightarrow x = y)$
- \leq , $=$, $<$ are antisymmetric but $|$ is not
- transitive iff $\forall x, y, z \in A (((x, y) \in R \wedge (y, z) \in R) \rightarrow (x, z) \in R)$
- \leq , $=$, $<$, and $|$ are transitive

R^* is the reflexive and transitive closure of R

Equivalence relations

Definition

A relation R on a set A is an equivalence relation iff it is reflexive, symmetric and transitive

Equivalence relations

Definition

A relation R on a set A is an equivalence relation iff it is reflexive, symmetric and transitive

- Let Σ^* be the set of strings over alphabet Σ . The relation $\{(s, t) \in \Sigma^* \times \Sigma^* \mid |s| = |t|\}$ is an equivalence relation

Equivalence relations

Definition

A relation R on a set A is an equivalence relation iff it is reflexive, symmetric and transitive

- Let Σ^* be the set of strings over alphabet Σ . The relation $\{(s, t) \in \Sigma^* \times \Sigma^* \mid |s| = |t|\}$ is an equivalence relation
- $|$ on integers is not an equivalence relation.

Equivalence relations

Definition

A relation R on a set A is an equivalence relation iff it is reflexive, symmetric and transitive

- Let Σ^* be the set of strings over alphabet Σ . The relation $\{(s, t) \in \Sigma^* \times \Sigma^* \mid |s| = |t|\}$ is an equivalence relation
- $|$ on integers is not an equivalence relation.
- For integer $m > 1$ the relation $\equiv \pmod{m}$ is an equivalence relation on integers

Equivalence classes

Definition

Let R be an equivalence relation on a set A and $a \in A$. Let

$$[a]_R = \{s \mid (a, s) \in R\}$$

be the equivalence class of a w.r.t. R

Equivalence classes

Definition

Let R be an equivalence relation on a set A and $a \in A$. Let

$$[a]_R = \{s \mid (a, s) \in R\}$$

be the equivalence class of a w.r.t. R

If $b \in [a]_R$ then b is called a representative of the equivalence class

Theorem

Result

Let R be an equivalence relation on A and $a, b \in A$. The following three statements are equivalent

- 1 aRb
- 2 $[a]_R = [b]_R$
- 3 $[a]_R \cap [b]_R \neq \emptyset$

Theorem

Result

Let R be an equivalence relation on A and $a, b \in A$. The following three statements are equivalent

- 1 aRb
- 2 $[a]_R = [b]_R$
- 3 $[a]_R \cap [b]_R \neq \emptyset$

Proof in book

Partitions of a set

Definition

A partition of a set A is a collection of disjoint, nonempty subsets that have A as their union. In other words, the collection of subsets $A_i \subseteq A$ with $i \in I$ (where I is an index set) forms a partition of A iff

- 1 $A_i \neq \emptyset$ for all $i \in I$
- 2 $A_i \cap A_j = \emptyset$ for all $i \neq j \in I$
- 3 $\bigcup_{i \in I} A_i = A$

Result

Theorem

- 1 If R is an equivalence on A , then the equivalence classes of R form a partition of A
- 2 Conversely, given a partition $\{A_i \mid i \in I\}$ of A there exists an equivalence relation R that has exactly the sets $A_i, i \in I$, as its equivalence classes

Result

Theorem

- 1 If R is an equivalence on A , then the equivalence classes of R form a partition of A
- 2 Conversely, given a partition $\{A_i \mid i \in I\}$ of A there exists an equivalence relation R that has exactly the sets $A_i, i \in I$, as its equivalence classes

Proof in book

Sequences

Sequences are ordered lists of elements

2, 3, 5, 7, 11, 13, 17, 19, ... or a, b, c, d, \dots, y, z

Sequences

Sequences are ordered lists of elements

2, 3, 5, 7, 11, 13, 17, 19, ... or a, b, c, d, \dots, y, z

Definition

A sequence over a set S is a function f from a subset of the integers (typically \mathbb{N} or \mathbb{Z}^+) to the set S . If the domain of f is finite then the sequence is finite

Examples

$f : \mathbb{Z}^+ \rightarrow \mathbb{Q}$ is $f(n) = 1/n$ defines the sequence

Examples

$f : \mathbb{Z}^+ \rightarrow \mathbb{Q}$ is $f(n) = 1/n$ defines the sequence

$$1, 1/2, 1/3, 1/4, \dots$$

Examples

$f : \mathbb{Z}^+ \rightarrow \mathbb{Q}$ is $f(n) = 1/n$ defines the sequence

$$1, 1/2, 1/3, 1/4, \dots$$

Assuming $a_n = f(n)$, the sequence is also written a_1, a_2, a_3, \dots

or as $\{a_n\}_{n \in \mathbb{Z}^+}$

Examples

$f : \mathbb{Z}^+ \rightarrow \mathbb{Q}$ is $f(n) = 1/n$ defines the sequence

$$1, 1/2, 1/3, 1/4, \dots$$

Assuming $a_n = f(n)$, the sequence is also written a_1, a_2, a_3, \dots

or as $\{a_n\}_{n \in \mathbb{Z}^+}$

$g : \mathbb{N} \rightarrow \mathbb{N}$ is $g(n) = n^2$ defines the sequence

$$0, 1, 4, 9, \dots$$

Examples

$f : \mathbb{Z}^+ \rightarrow \mathbb{Q}$ is $f(n) = 1/n$ defines the sequence

$$1, 1/2, 1/3, 1/4, \dots$$

Assuming $a_n = f(n)$, the sequence is also written a_1, a_2, a_3, \dots

or as $\{a_n\}_{n \in \mathbb{Z}^+}$

$g : \mathbb{N} \rightarrow \mathbb{N}$ is $g(n) = n^2$ defines the sequence

$$0, 1, 4, 9, \dots$$

Assuming $b_n = g(n)$, also written b_0, b_1, b_2, \dots or as $\{b_n\}_{n \in \mathbb{N}}$

Geometric and arithmetic progressions

- A geometric progression is a sequence of the form

$$a, ar, ar^2, ar^3, \dots, ar^n, \dots$$

Geometric and arithmetic progressions

- A geometric progression is a sequence of the form

$$a, ar, ar^2, ar^3, \dots, ar^n, \dots$$

- Example $\{b_n\}_{n \in \mathbb{N}}$ with $b_n = 6(1/3)^n$

Geometric and arithmetic progressions

- A geometric progression is a sequence of the form

$$a, ar, ar^2, ar^3, \dots, ar^n, \dots$$

- Example $\{b_n\}_{n \in \mathbb{N}}$ with $b_n = 6(1/3)^n$

- An arithmetic progression is a sequence of the form

$$a, a + d, a + 2d, a + 3d, \dots, a + nd, \dots$$

Geometric and arithmetic progressions

- A geometric progression is a sequence of the form

$$a, ar, ar^2, ar^3, \dots, ar^n, \dots$$

- Example $\{b_n\}_{n \in \mathbb{N}}$ with $b_n = 6(1/3)^n$

- An arithmetic progression is a sequence of the form

$$a, a + d, a + 2d, a + 3d, \dots, a + nd, \dots$$

- Example $\{c_n\}_{n \in \mathbb{N}}$ with $c_n = 7 - 3n$

Geometric and arithmetic progressions

- A geometric progression is a sequence of the form

$$a, ar, ar^2, ar^3, \dots, ar^n, \dots$$

- Example $\{b_n\}_{n \in \mathbb{N}}$ with $b_n = 6(1/3)^n$

- An arithmetic progression is a sequence of the form

$$a, a + d, a + 2d, a + 3d, \dots, a + nd, \dots$$

- Example $\{c_n\}_{n \in \mathbb{N}}$ with $c_n = 7 - 3n$

where the initial elements a , the common ratio r and the common difference d are real numbers

Recurrence relations

Definition

A recurrence relation for $\{a_n\}_{n \in \mathbb{N}}$ is an equation that expresses a_n in terms of one or more of the elements a_0, a_1, \dots, a_{n-1}

Recurrence relations

Definition

A recurrence relation for $\{a_n\}_{n \in \mathbb{N}}$ is an equation that expresses a_n in terms of one or more of the elements a_0, a_1, \dots, a_{n-1}

- Typically the recurrence relation expresses a_n in terms of just a fixed number of previous elements (such as $a_n = g(a_{n-1}, a_{n-2})$)

Recurrence relations

Definition

A recurrence relation for $\{a_n\}_{n \in \mathbb{N}}$ is an equation that expresses a_n in terms of one or more of the elements a_0, a_1, \dots, a_{n-1}

- Typically the recurrence relation expresses a_n in terms of just a fixed number of previous elements (such as $a_n = g(a_{n-1}, a_{n-2})$)
- The initial conditions specify the first elements of the sequence, before the recurrence relation applies

Recurrence relations

Definition

A recurrence relation for $\{a_n\}_{n \in \mathbb{N}}$ is an equation that expresses a_n in terms of one or more of the elements a_0, a_1, \dots, a_{n-1}

- Typically the recurrence relation expresses a_n in terms of just a fixed number of previous elements (such as $a_n = g(a_{n-1}, a_{n-2})$)
- The initial conditions specify the first elements of the sequence, before the recurrence relation applies
- A sequence is called a solution of a recurrence relation iff its terms satisfy the recurrence relation

Rabbits and Fibonacci sequence

After 1 month, a pair of rabbits is placed on an island

Rabbits and Fibonacci sequence

After 1 month, a pair of rabbits is placed on an island

After every 2 months, each pair of rabbits produces a new pair of rabbits

Rabbits and Fibonacci sequence

After 1 month, a pair of rabbits is placed on an island

After every 2 months, each pair of rabbits produces a new pair of rabbits

Find a recurrence relation for number of pairs of rabbits after $n \in \mathbb{Z}^+$ months assuming no rabbits die

Rabbits and Fibonacci sequence

After 1 month, a pair of rabbits is placed on an island

After every 2 months, each pair of rabbits produces a new pair of rabbits

Find a recurrence relation for number of pairs of rabbits after $n \in \mathbb{Z}^+$ months assuming no rabbits die

Answer is the Fibonacci sequence

$$\begin{cases} f(1) = 1 \\ f(2) = 1 \\ f(n) = f(n-1) + f(n-2) \text{ for } n > 2 \end{cases}$$

Rabbits and Fibonacci sequence

After 1 month, a pair of rabbits is placed on an island

After every 2 months, each pair of rabbits produces a new pair of rabbits

Find a recurrence relation for number of pairs of rabbits after $n \in \mathbb{Z}^+$ months assuming no rabbits die

Answer is the Fibonacci sequence

$$\begin{cases} f(1) = 1 \\ f(2) = 1 \\ f(n) = f(n-1) + f(n-2) \text{ for } n > 2 \end{cases}$$

Yields the sequence 1, 1, 2, 3, 5, 8, 13, ...

Solving recurrence relations

- Finding a formula for the n^{th} term of the sequence generated by a recurrence relation is called solving the recurrence relation

Solving recurrence relations

- Finding a formula for the n^{th} term of the sequence generated by a recurrence relation is called solving the recurrence relation
- Such a formula is called a closed formula

Solving recurrence relations

- Finding a formula for the n^{th} term of the sequence generated by a recurrence relation is called solving the recurrence relation
- Such a formula is called a closed formula
- Various more advanced methods for solving recurrence relations are covered in Chapter 8 of the book (not part of this course)

Solving recurrence relations

- Finding a formula for the n^{th} term of the sequence generated by a recurrence relation is called solving the recurrence relation
- Such a formula is called a closed formula
- Various more advanced methods for solving recurrence relations are covered in Chapter 8 of the book (not part of this course)
- Here we illustrate by example the method of iteration in which we need to guess the formula

Solving recurrence relations

- Finding a formula for the n^{th} term of the sequence generated by a recurrence relation is called solving the recurrence relation
- Such a formula is called a closed formula
- Various more advanced methods for solving recurrence relations are covered in Chapter 8 of the book (not part of this course)
- Here we illustrate by example the method of iteration in which we need to guess the formula
- The guess can be proved correct by the method of induction (to be covered)

Iterative solution - working upwards

Forward substitution

Iterative solution - working upwards

Forward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

Iterative solution - working upwards

Forward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$a_2 = 2 + 3$$

Iterative solution - working upwards

Forward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$a_2 = 2 + 3$$

$$a_3 = (2 + 3) + 3 = 2 + 3 \cdot 2$$

Iterative solution - working upwards

Forward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$a_2 = 2 + 3$$

$$a_3 = (2 + 3) + 3 = 2 + 3 \cdot 2$$

$$a_4 = (2 + 2 \cdot 3) + 3 = 2 + 3 \cdot 3$$

Iterative solution - working upwards

Forward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$a_2 = 2 + 3$$

$$a_3 = (2 + 3) + 3 = 2 + 3 \cdot 2$$

$$a_4 = (2 + 2 \cdot 3) + 3 = 2 + 3 \cdot 3$$

\vdots

$$a_n = a_{n-1} + 3 = (2 + 3 \cdot (n - 2)) + 3 = 2 + 3 \cdot (n - 1)$$

Iterative solution - working downward

Backward substitution

Iterative solution - working downward

Backward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$a_n = a_{n-1} + 3$$

Iterative solution - working downward

Backward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$\begin{aligned} a_n &= a_{n-1} + 3 \\ &= (a_{n-2} + 3) + 3 = a_{n-2} + 3 \cdot 2 \end{aligned}$$

Iterative solution - working downward

Backward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$\begin{aligned} a_n &= a_{n-1} + 3 \\ &= (a_{n-2} + 3) + 3 = a_{n-2} + 3 \cdot 2 \\ &= (a_{n-3} + 3) + 3 \cdot 2 = a_{n-3} + 3 \cdot 3 \end{aligned}$$

Iterative solution - working downward

Backward substitution

$$a_n = a_{n-1} + 3 \text{ for } n \geq 2 \text{ with } a_1 = 2$$

$$\begin{aligned} a_n &= a_{n-1} + 3 \\ &= (a_{n-2} + 3) + 3 = a_{n-2} + 3 \cdot 2 \\ &= (a_{n-3} + 3) + 3 \cdot 2 = a_{n-3} + 3 \cdot 3 \\ &\quad \vdots \\ &= a_2 + 3(n-2) = (a_1 + 3) + 3 \cdot (n-2) = 2 + 3 \cdot (n-1) \end{aligned}$$

Compound interest

- Suppose a person deposits £1000 in a savings account yielding 3% per year with interest compounded annually. How much is in the account after 20 years?

Compound interest

- Suppose a person deposits £1000 in a savings account yielding 3% per year with interest compounded annually. How much is in the account after 20 years?
- Let P_n denote amount after n years

Compound interest

- Suppose a person deposits £1000 in a savings account yielding 3% per year with interest compounded annually. How much is in the account after 20 years?
- Let P_n denote amount after n years
- $P_n = P_{n-1} + 0.03 P_{n-1} = (1.03)P_{n-1}$

Compound interest

- Suppose a person deposits £1000 in a savings account yielding 3% per year with interest compounded annually. How much is in the account after 20 years?
- Let P_n denote amount after n years
- $P_n = P_{n-1} + 0.03 P_{n-1} = (1.03)P_{n-1}$
- The initial condition $P_0 = 1000$.

Compound interest

- Suppose a person deposits £1000 in a savings account yielding 3% per year with interest compounded annually. How much is in the account after 20 years?
- Let P_n denote amount after n years
- $P_n = P_{n-1} + 0.03 P_{n-1} = (1.03)P_{n-1}$
- The initial condition $P_0 = 1000$.
- $P_1 = (1.03) P_0, \dots, P_n = (1.03)P_{n-1} = (1.03)^n P_0$

Compound interest

- Suppose a person deposits £1000 in a savings account yielding 3% per year with interest compounded annually. How much is in the account after 20 years?
- Let P_n denote amount after n years
- $P_n = P_{n-1} + 0.03 P_{n-1} = (1.03)P_{n-1}$
- The initial condition $P_0 = 1000$.
- $P_1 = (1.03) P_0, \dots, P_n = (1.03)P_{n-1} = (1.03)^n P_0$
- $P_{20} = (1.03)^{20} 1000 = 1,806$

Common sequences

TABLE 1 Some Useful Sequences.

<i>n</i> th Term	First 10 Terms
n^2	1, 4, 9, 16, 25, 36, 49, 64, 81, 100, ...
n^3	1, 8, 27, 64, 125, 216, 343, 512, 729, 1000, ...
n^4	1, 16, 81, 256, 625, 1296, 2401, 4096, 6561, 10000, ...
2^n	2, 4, 8, 16, 32, 64, 128, 256, 512, 1024, ...
3^n	3, 9, 27, 81, 243, 729, 2187, 6561, 19683, 59049, ...
$n!$	1, 2, 6, 24, 120, 720, 5040, 40320, 362880, 3628800, ...
f_n	1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, ...

Summations

Given a sequence $\{a_n\}$, the sum of terms $a_m, a_{m+1}, \dots, a_\ell$ is

$$a_m + a_{m+1} + \dots + a_\ell$$

Summations

Given a sequence $\{a_n\}$, the sum of terms $a_m, a_{m+1}, \dots, a_\ell$ is

$$a_m + a_{m+1} + \dots + a_\ell$$

$$\sum_{j=m}^{\ell} a_j \quad \text{or} \quad \sum_{m \leq j \leq \ell} a_j$$

Summations

Given a sequence $\{a_n\}$, the sum of terms $a_m, a_{m+1}, \dots, a_\ell$ is

$$a_m + a_{m+1} + \dots + a_\ell$$

$$\sum_{j=m}^{\ell} a_j \quad \text{or} \quad \sum_{m \leq j \leq \ell} a_j$$

The variable j is called the index of summation

More generally for an index set S

$$\sum_{j \in S} a_j$$

Useful summation formulas

TABLE 2 Some Useful Summation Formulae.

<i>Sum</i>	<i>Closed Form</i>
$\sum_{k=0}^n ar^k \quad (r \neq 0)$	$\frac{ar^{n+1} - a}{r - 1}, r \neq 1$
$\sum_{k=1}^n k$	$\frac{n(n+1)}{2}$
$\sum_{k=1}^n k^2$	$\frac{n(n+1)(2n+1)}{6}$
$\sum_{k=1}^n k^3$	$\frac{n^2(n+1)^2}{4}$
$\sum_{k=0}^{\infty} x^k, x < 1$	$\frac{1}{1-x}$
$\sum_{k=1}^{\infty} kx^{k-1}, x < 1$	$\frac{1}{(1-x)^2}$

Products

Given a sequence $\{a_n\}$, the product of terms $a_m, a_{m+1}, \dots, a_\ell$ is

$$a_m \cdot a_{m+1} \cdot \dots \cdot a_\ell$$

$$\prod_{j=m}^{\ell} a_j \quad \text{or} \quad \prod_{m \leq j \leq \ell} a_j$$

More generally for a finite index set S one writes

$$\prod_{j \in S} a_j$$