

## Overview

# Elements of Programming Languages

## Lecture 13: Small-step semantics and type safety

James Cheney

University of Edinburgh

November 8, 2016

- For the remaining lectures we consider some *cross-cutting* considerations for programming language design.
  - Last time: Imperative programming
- Today:
  - Finer-grained (small-step) evaluation
  - Type safety

## Refresher

- In the first 6 lectures we covered:
  - Basic arithmetic ( $L_{\text{Arith}}$ )
  - Conditionals and booleans ( $L_{\text{If}}$ )
  - Variables and let-binding ( $L_{\text{Let}}$ )
  - Functions and recursion ( $L_{\text{Rec}}$ )
  - Data structures ( $L_{\text{Data}}$ )
- formalized using big-step evaluation ( $e \Downarrow v$ ) and type judgments ( $\Gamma \vdash e : \tau$ )
- and implemented using Scala interpreters (CW1)

## Limitations of big-step semantics

- Big-step semantics is convenient, but also limited
- It says how to evaluate the “whole program” (expression) to its “final value”
- *But what if there is no final value?*
  - Expressions like  $1 + \text{true}$  simply don't evaluate
  - Nonterminating programs don't evaluate either, but for a different reason!
- As we will see in later lectures, it is also difficult to deal with other features, like exceptions, using big-step semantics

## Small-step semantics

- We will now consider an alternative: *small-step semantics*

$$e \mapsto e'$$

- which says how to evaluate an expression “one step at a time”
- If  $e_0 \mapsto \dots \mapsto e_n$  then we write  $e_0 \mapsto^* e_n$ . (in particular, for  $n = 0$  we have  $e_0 \mapsto^* e_0$ )
- We want it to be the case that  $e \mapsto^* v$  if and only if  $e \Downarrow v$ .
- But  $\mapsto$  provides more detail about how this happens.
- It also allows expressions to “go wrong” (get stuck before reaching a value)

Small-step semantics:  $L_{If}$ 
 $e \mapsto e'$  for  $L_{If}$ 

$$\frac{}{v == v \mapsto \text{true}} \quad \frac{v_1 \neq v_2}{v_1 == v_2 \mapsto \text{false}}$$

$$\frac{e \mapsto e'}{\text{if } e \text{ then } e_1 \text{ else } e_2 \mapsto \text{if } e' \text{ then } e_1 \text{ else } e_2}$$

$$\frac{}{\text{if true then } e_1 \text{ else } e_2 \mapsto e_1}$$

$$\frac{}{\text{if false then } e_1 \text{ else } e_2 \mapsto e_2}$$

- If the conditional test is not a value, evaluate it one step
  - Otherwise, evaluate the corresponding branch
- $$\text{if } 1 == 2 \text{ then } 3 \text{ else } 4 \mapsto \text{if false then } 3 \text{ else } 4 \mapsto 4$$

Small-step semantics:  $L_{Arith}$ 
 $e \mapsto e'$  for  $L_{Arith}$ 

$$\frac{e_1 \mapsto e'_1}{e_1 \oplus e_2 \mapsto e'_1 \oplus e_2} \quad \frac{e_2 \mapsto e'_2}{v_1 \oplus e_2 \mapsto v_1 \oplus e'_2}$$

$$\frac{}{v_1 + v_2 \mapsto v_1 +_{\mathbb{N}} v_2} \quad \frac{}{v_1 \times v_2 \mapsto v_1 \times_{\mathbb{N}} v_2}$$

- If the first subexpression of  $\oplus$  can take a step, apply it
- If the first subexpression is a value and the second can take a step, apply it
- If both sides are values, perform the operation
- Example:

$$1 + (2 \times 3) \mapsto 1 + 6 \mapsto 7$$

Small-step semantics:  $L_{Let}$ 
 $e \mapsto e'$  for  $L_{Let}$ 

$$\frac{e_1 \mapsto e'_1}{\text{let } x = e_1 \text{ in } e_2 \mapsto \text{let } x = e'_1 \text{ in } e_2}$$

$$\frac{}{\text{let } x = v_1 \text{ in } e_2 \mapsto e_2[v_1/x]}$$

- If the expression  $e_1$  is not yet a value, evaluate it one step
- Otherwise, substitute it and proceed
- Example:

$$\text{let } x = 1 + 1 \text{ in } x \times x \mapsto \text{let } x = 2 \text{ in } x \times x \mapsto 2 \times 2 \mapsto 4$$



Small-step semantics:  $L_{Lam}$  $e \mapsto e'$  for  $L_{Lam}$ 

$$\frac{e_1 \mapsto e'_1}{e_1 e_2 \mapsto e'_1 e_2} \quad \frac{e_2 \mapsto e'_2}{v_1 e_2 \mapsto v_1 e'_2}$$

$$\overline{(\lambda x. e) v \mapsto e[v/x]}$$

- If the function part is not a value, evaluate it one step
- If the function is a value and the argument isn't, evaluate it one step
- If both function and argument are values, substitute and proceed

$$\begin{aligned} ((\lambda x. \lambda y. x + y) 1) 2 &\mapsto (\lambda y. 1 + y) 2 \\ &\mapsto 1 + 2 \mapsto 3 \end{aligned}$$

Small-step semantics:  $L_{Rec}$  $e \mapsto e'$  for  $L_{Rec}$ 

$$\overline{(\text{rec } f(x). e) v \mapsto e[\text{rec } f(x). e/f, v/x]}$$

- Same rules for evaluation inside application
- Note that we need to substitute  $\text{rec } f(x). e$  for  $f$ .
- Suppose *fact* is the factorial function:

$$\begin{aligned} \text{fact } 2 &\mapsto \text{if } 2 == 0 \text{ then } 1 \text{ else } 2 \times \text{fact}(2 - 1) \\ &\mapsto \text{if false then } 1 \text{ else } 2 \times \text{fact}(2 - 1) \\ &\mapsto 2 \times \text{fact}(2 - 1) \mapsto 2 \times \text{fact}(1) \\ &\mapsto 2 \times (\text{if } 1 == 0 \text{ then } 1 \text{ else } 1 \times \text{fact}(1 - 1)) \\ &\mapsto 2 \times (\text{if false then } 1 \text{ else } 1 \times \text{fact}(1 - 1)) \\ &\mapsto 2 \times (1 \times \text{fact}(1 - 1)) \mapsto 2 \times (1 \times \text{fact}(0)) \\ &\mapsto^* 2 \times (1 \times 1) \mapsto 2 \times 1 \mapsto 2 \end{aligned}$$



## Judgments and Rules, in general

- A *judgment* is a relation among one or more abstract syntax trees.
- Examples so far:  $e \Downarrow v$ ,  $\Gamma \vdash e : \tau$ ,  $e \mapsto e'$
- We have been defining judgments using *rules* of the form:

$$\overline{Q} \quad \frac{P_1 \quad \dots \quad P_n}{Q}$$

- where  $P_1, \dots, P_n$  and  $Q$  are judgments.



## Meaning of Rules

- A rule of the form:

$$\overline{Q}$$

is called an *axiom*. It says that  $Q$  is always derivable.

- A rule of the form

$$\frac{P_1 \quad \dots \quad P_n}{Q}$$

says that judgment  $Q$  is derivable if  $P_1, \dots, P_n$  are derivable.

- Symbols like  $e, v, \tau$  in rules stand for arbitrary expressions, values, or types.
- (If you have taken Logic Programming: These rules are a lot like Prolog clauses!)



## Rule induction

### Induction on derivations of $e \Downarrow v$

Suppose  $P(-, -)$  is a predicate over pairs of expressions and values. If:

- $P(v, v)$  holds for all values  $v$
- If  $P(e_1, v_1)$  and  $P(e_2, v_2)$  then  $P(e_1 + e_2, v_1 +_{\mathbb{N}} v_2)$
- If  $P(e_1, v_1)$  and  $P(e_2, v_2)$  then  $P(e_1 \times e_2, v_1 \times_{\mathbb{N}} v_2)$

then  $e \Downarrow v$  implies  $P(e, v)$ .

- Rule induction can be derived from mathematical induction on the size (or height) of the derivation tree.
- (Much like structural induction.)
- We won't formally prove this.

## Example: $e \Downarrow v$ implies $e \mapsto^* v$

- As an example, we'll show a few cases of the forward direction of:

### Theorem (Equivalence of big-step and small-step evaluation)

$e \Downarrow v$  if and only if  $e \mapsto^* v$ .

### Base case.

If the derivation is of the form

$$\overline{n \Downarrow n}$$

for some number  $n$ , then  $e = n$  is already a value  $v = n$ , so no steps are needed to evaluate it, i.e.  $n \mapsto^* n$  in zero steps.  $\square$

## Example: $e \Downarrow v$ implies $e \mapsto^* v$

### Inductive case.

If the derivation is of the form

$$\frac{e_1 \Downarrow v_2 \quad e_2 \Downarrow v_2}{e_1 + e_2 \Downarrow v_1 +_{\mathbb{N}} v_2}$$

then by induction, we know  $e_1 \mapsto^* v_1$  and  $e_2 \mapsto^* v_2$ . Using the small-step rules, we can then show

$$e_1 + e_2 \mapsto^* v_1 + e_2 \mapsto^* v_1 + v_2 \mapsto v_1 +_{\mathbb{N}} v_2$$

$\square$

- The case for  $\times$  is similar.

## Type soundness

- The central property of a type system is *soundness*.
- Roughly speaking, soundness means “well-typed programs don't go wrong” [Milner].
- But what exactly does “go wrong” mean?
  - For large-step: hard to say
  - For small-step: “go wrong” means “stuck” expression  $e$  that is not a value and cannot take a step.
- We could show something like:

### Theorem (Soundness)

If  $\vdash e : \tau$  and  $e \mapsto^* v$  then  $\vdash v : \tau$ .

- This says that if an expression evaluates to a value, then the value has the right type.

## Type soundness revisited

- We can decompose soundness into two parts:

### Lemma (Progress)

If  $\vdash e : \tau$  then either  $e$  is a value or for some  $e'$  we have  $e \mapsto e'$ .

### Lemma (Preservation)

If  $\vdash e : \tau$  and  $e \mapsto e'$  then  $\vdash e' : \tau$

- Combining these two, can show:

### Theorem (Soundness)

If  $\vdash e : \tau$  and  $e \mapsto^* v$  then  $\vdash v : \tau$ .

- We will *sketch* these properties for  $L_{lf}$  (leaving out a lot of formal detail)



## Progress for $L_{lf}$

Progress is proved by induction on  $\vdash e : \tau$  derivations. We show some representative cases.

### Progress for $+$ .

$$\frac{\vdash e_1 : \text{int} \quad e_2 : \text{int}}{\vdash e_1 + e_2 : \text{int}}$$

If the derivation is of the above form, then by induction  $e_1$  is either a value or can take a step, and likewise for  $e_2$ . There are three cases.

- If  $e_1 \mapsto e'_1$  then  $e_1 + e_2 \mapsto e'_1 + e_2$ .
- If  $e_1$  is a value  $v_1$  and  $e_2 \mapsto e'_2$ , then  $v_1 + e_2 \mapsto v_1 + e'_2$ .
- If both  $e_1$  and  $e_2$  are values then they must both be numbers  $n_1, n_2 \in \mathbb{N}$ , so  $e_1 + e_2 \mapsto n_1 +_{\mathbb{N}} n_2$ .



## Progress for $L_{lf}$

### Progress for *if*.

If the derivation is of the form

$$\frac{\vdash e : \text{bool} \quad \vdash e_1 : \tau \quad \vdash e_2 : \tau}{\vdash \text{if } e \text{ then } e_1 \text{ else } e_2 : \tau}$$

then by induction, either  $e$  is a value or can take a step. There are two cases:

- If  $e \mapsto e'$  then  
if  $e$  then  $e_1$  else  $e_2 \mapsto \text{if } e' \text{ then } e_1 \text{ else } e_2$ .
- If  $e$  is a value, it must be either true or false. Then  
if true then  $e_1$  else  $e_2 \mapsto e_1$  or  
if false then  $e_1$  else  $e_2 \mapsto e_2$ .



## Preservation for $L_{lf}$

Preservation is proved by induction on the structure of  $\vdash e : \tau$ . We'll consider some representative cases:

### Preservation for $+$ .

$$\frac{\vdash e_1 : \text{int} \quad \vdash e_2 : \text{int}}{\vdash e_1 + e_2 : \text{int}}$$

If the derivation is of the above form, there are three cases.

- If  $e_i = v_i$  and  $v_1 + v_2 \mapsto v_1 +_{\mathbb{N}} v_2$  then obviously  $\vdash v_1 +_{\mathbb{N}} v_2 : \text{int}$ .
- If  $e_1 + e_2 \mapsto e'_1 + e_2$  where  $e_1 \mapsto e'_1$ , then since  $\vdash e_1 : \text{int}$ , we have  $\vdash e'_1 : \text{int}$ , so  $\vdash e'_1 + e_2 : \text{int}$  also.
- The case where  $e_1 = v_1$  and  $v_1 + e_2 \mapsto v_1 + e'_2$  is similar.



## Preservation for $L_{If}$

### Preservation for if.

If the derivation is of the form

$$\frac{\vdash e : \text{bool} \quad \vdash e_1 : \tau \quad \vdash e_2 : \tau}{\vdash \text{if } e \text{ then } e_1 \text{ else } e_2 : \tau}$$

then there are three cases:

- If  $\text{if } e \text{ then } e_1 \text{ else } e_2 \mapsto \text{if } e' \text{ then } e_1 \text{ else } e_2$  where  $e \mapsto e'$ , then by induction we can show that  $\vdash e' : \text{bool}$  and  $\vdash \text{if } e' \text{ then } e_1 \text{ else } e_2 : \tau$ .
- If  $e = \text{true}$  then  $\text{if true then } e_1 \text{ else } e_2 \mapsto e_1$ , so we already know  $\vdash e_1 : \tau$ .
- The case for  $\text{if false then } e_1 \text{ else } e_2 \mapsto e_2$  is similar.



## Type soundness for $L_{Let}$ [non-examinable]

- Progress: straightforward (a “let” can always take a step)
- Preservation: Suppose we have

$$\frac{\vdash v_1 : \tau' \quad x:\tau' \vdash e_2 : \tau}{\vdash \text{let } x = v_1 \text{ in } e_2 : \tau} \quad \frac{}{\text{let } x = v_1 \text{ in } e_2 \mapsto e_2[v_1/x]}$$

We need to show that  $\vdash e_2[v_1/x] : \tau$

- For this we need a *substitution lemma*

### Lemma (Substitution)

If  $\Gamma, x:\tau' \vdash e : \tau$  and  $\Gamma \vdash e' : \tau'$  then  $\Gamma \vdash e[e'/x] : \tau$

## Type soundness for $L_{Rec}$ [non-examinable]

- Progress: If an application term is well-formed:

$$\frac{\vdash e_1 : \tau_1 \rightarrow \tau_2 \quad \vdash e_2 : \tau_1}{\vdash e_1 e_2 : \tau_2}$$

then by induction,  $e_1$  is either a value or  $e_1 \mapsto e'_1$  for some  $e'_1$ . If it is a value, it must be either a lambda-expression or a recursive function, so  $e_1 e_2$  can take a step.

Otherwise,  $e_1 e_2 \mapsto e'_1 e_2$ .

- Preservation: Similar to let, using substitution lemma for the cases

$$\begin{aligned} (\lambda x. e) v &\mapsto e[v/x] \\ (\text{rec } f(x). e) v &\mapsto e[\text{rec } f(x). e/f, v/x] \end{aligned}$$

## Summary

- Today we have presented
  - Small-step evaluation: a finer-grained semantics
  - Induction on derivations
  - Type soundness (details for  $L_{If}$ )
  - Sketch of type soundness for  $L_{Rec}$  [Non-examinable]
- Deep breath: No more proofs from now on.
- Remaining lectures cover cross-cutting language features, which often have subtle interactions with each other
  - Next time: Guest lecture by Michel Steuwer on *DSLs and rewrite-based optimizations for performance-portable parallel programming*