

# Formal Modeling in Cognitive Science

## Lecture 20: Joint, Marginal, and Conditional Distributions

Steve Renals (notes by Frank Keller)

School of Informatics  
University of Edinburgh  
s.renals@ed.ac.uk

26 February 2007

- 1 Distributions
  - Joint Distributions
  - Marginal Distributions
  - Conditional Distributions
  
- 2 Independence

# Joint Distributions

Previously, we introduced  $P(A \cap B)$ , the *probability of the intersection* of the two events  $A$  and  $B$ .

Let these events be described by the random variables  $X$  at value  $x$  and  $Y$  at value  $y$ . Then we can write:

$$P(A \cap B) = P(X = x \cap Y = y) = P(X = x, Y = y)$$

This is referred to as the *joint probability* of  $X = x$  and  $Y = y$ .

Note: often the term joint probability and the notation  $P(A, B)$  is also used for the probability of the intersection of two events.

# Joint Distributions

The notion of the joint probability can be generalized to distributions:

## Definition: Joint Probability Distribution

If  $X$  and  $Y$  are discrete random variables, the function given by  $f(x, y) = P(X = x, Y = y)$  for each pair of values  $(x, y)$  within the range of  $X$  is called the joint probability distribution of  $X$  and  $Y$ .

## Definition: Joint Cumulative Distribution

If  $X$  and  $Y$  are a discrete random variables, the function given by:

$$F(x, y) = P(X \leq x, Y \leq y) = \sum_{s \leq x} \sum_{t \leq y} f(s, t) \text{ for } -\infty < x, y < \infty$$

where  $f(s, t)$  is the value of the joint probability distribution of  $X$  and  $Y$  at  $(s, t)$ , is the joint cumulative distribution of  $X$  and  $Y$ .

## Example: Corpus Data

Assume you have a corpus of a 100 words (a corpus is a collection of text; see Informatics 1B). You tabulate the words, their frequencies and probabilities in the corpus:

$w$	$c(w)$	$P(w)$	$x$	$y$
the	30	0.30	3	1
to	18	0.18	2	1
will	16	0.16	4	1
of	10	0.10	2	1
Earth	7	0.07	5	2
on	6	0.06	2	1
probe	4	0.04	5	2
some	3	0.03	4	2
Comet	3	0.03	5	2
BBC	3	0.03	3	0

## Example: Corpus Data

We can now define the following random variables:

- $X$ : the length of the word;
- $Y$ : number of vowels in the word.

Examples for probability distributions:

- $f_X(5) = P(\text{Earth}) + P(\text{probe}) + P(\text{Comet}) = 0.14$ ;
- $f_Y(2) = P(\text{Earth}) + P(\text{probe}) + P(\text{some}) + P(\text{Comet}) = 0.17$ .

Examples for cumulative distributions:

- $F_X(3) = f_X(2) + f_X(3) = 0.34 + 0.33 = 0.67$ ;
- $F_Y(1) = f_X(0) + f_X(1) = 0.03 + 0.80 = 0.83$ .

## Example: Corpus Data

Now compute the joint distribution of  $X$  and  $Y$  as  $f(x, y) = P(X = x, Y = y)$ .

Examples:

- $f(2, 1) = P(\text{to}) + P(\text{of}) + P(\text{on}) = 0.18 + 0.10 + 0.06 = 0.34$ ;
- $f(3, 0) = P(\text{BBC}) = 0.03$ ;
- $f(4, 3) = 0$ .

Full distribution:

		x			
		2	3	4	5
y	0	0	0.03	0	0
	1	0.34	0.30	0.16	0
	2	0	0	0.03	0.14

# Marginal Distributions

If we 'project' one of the two dimensions of a joint distributions, we obtain a marginal distributions:

## Definition: Marginal Distribution

If  $X$  and  $Y$  are discrete random variables and  $f(x, y)$  is the value of their joint probability distribution at  $(x, y)$ , the functions given by:

$$g(x) = \sum_y f(x, y) \quad \text{and} \quad h(y) = \sum_x f(x, y)$$

are the marginal distributions of  $X$  and  $Y$ , respectively.



## Example: Corpus Data

We had defined the following random variables:

- $X$ : the length of the word;
- $Y$ : number of vowels in the word.

Joint distribution of  $X$  and  $Y$ :

		x				$\sum_x f(x, y)$
		2	3	4	5	
y	0	0	0.03	0	0	0.03
	1	0.34	0.30	0.16	0	0.80
	2	0	0	0.03	0.14	0.17
$\sum_y f(x, y)$		0.34	0.33	0.19	0.14	

*Marginal distribution of  $Y$ . Marginal distribution of  $X$ .*

# Conditional Distributions

Previously, we defined the *conditional probability* of two events  $A$  and  $B$  as follows:

$$P(B|A) = \frac{P(A \cap B)}{P(A)}$$

Let these events be described by the random variable  $X = x$  and  $Y = y$ . Then we can write:

$$P(X = x|Y = y) = \frac{P(X = x, Y = y)}{P(Y = y)} = \frac{f(x, y)}{h(y)}$$

where  $f(x, y)$  is the joint probability distribution of  $X$  and  $Y$  and  $h(y)$  is the marginal marginal distribution of  $y$ .

# Conditional Distributions

## Definition: Conditional Distribution

If  $f(x, y)$  is the value of the joint probability distribution of the discrete random variables  $X$  and  $Y$  at  $(x, y)$  and  $h(y)$  is the value of the marginal distributions of  $Y$  at  $y$ , and  $g(x)$  is the value of the marginal distributions of  $X$  at  $x$ , then:

$$f(x|y) = \frac{f(x, y)}{h(y)} \quad \text{and} \quad w(y|x) = \frac{f(x, y)}{g(x)}$$

are the conditional distributions of  $X$  given  $Y = y$ , and of  $Y$  given  $X = x$ , respectively (for  $h(y) \neq 0$  and  $g(x) \neq 0$ ).

## Example: Corpus Data

Based on the joint distribution  $f(x, y)$  and the marginal distributions  $h(y)$  and  $g(x)$  from the previous example, we can compute the conditional distributions of  $X$  given  $Y = 1$ :

		x			
		2	3	4	5
y	1	$\frac{f(2,1)}{h(1)} = \frac{0.34}{0.80} = 0.43$	$\frac{f(3,1)}{h(1)} = \frac{0.30}{0.80} = 0.38$	$\frac{f(4,1)}{h(1)} = \frac{0.16}{0.80} = 0.20$	$\frac{f(5,1)}{h(1)} = \frac{0}{0.80} = 0$

# Independence

The notion of *independence* of events can also be generalized to probability distributions:

## Definition: Independence

If  $f(x, y)$  is the value of the joint probability distribution of the discrete random variables  $X$  and  $Y$  at  $(x, y)$ , and  $g(x)$  and  $h(y)$  are the values of the marginal distributions of  $X$  at  $x$  and  $Y$  at  $y$ , respectively, then  $X$  and  $Y$  are independent iff:

$$f(x, y) = g(x)h(y)$$

for all  $(x, y)$  within their range.

## Example: Corpus Data

Marginal distributions from the previous example:

		x				h(y)
		2	3	4	5	
y	0	0	0.03	0	0	0.03
	1	0.34	0.30	0.16	0	0.80
	2	0	0	0.03	0.14	0.17
g(x)		0.34	0.33	0.19	0.14	

Now compute  $g(x)h(y)$  for each cell in the table:

		x			
		2	3	4	5
y	0	0.01	0.01	0.01	0.00
	1	0.27	0.26	0.15	0.12
	2	0.06	0.06	0.03	0.02

X and Y are  
*not independent.*

# Summary

- A joint probability distribution returns a probability for each pair of values of two random variables.
- marginal distributions project one of the dimensions of a joint probability distribution;
- the conditional distribution is the joint distribution divided by the marginal distribution;
- two distributions are independent if the joint distribution is the same as the product of the two marginal distributions.