

The Matrix Cookbook

Kaare Brandt Petersen
Michael Syskind Pedersen

VERSION: JANUARY 5, 2005

What is this? These pages are a collection of facts (identities, approximations, inequalities, relations, ...) about matrices and matters relating to them. It is collected in this form for the convenience of anyone who wants a quick desktop reference .

Disclaimer: The identities, approximations and relations presented here were obviously not invented but collected, borrowed and copied from a large amount of sources. These sources include similar but shorter notes found on the internet and appendices in books - see the references for a full list.

Errors: Very likely there are errors, typos, and mistakes for which we apologize and would be grateful to receive corrections at kbp@imm.dtu.dk.

Its ongoing: The project of keeping a large repository of relations involving matrices is naturally ongoing and the version will be apparent from the date in the header.

Suggestions: Your suggestion for additional content or elaboration of some topics is most welcome at kbp@imm.dtu.dk.

Acknowledgements: We would like to thank the following for discussions, proofreading, extensive corrections and suggestions: Esben Hoegh-Rasmussen and Vasile Sima.

Keywords: Matrix algebra, matrix relations, matrix identities, derivative of determinant, derivative of inverse matrix, differentiate a matrix.

Contents

1	Basics	5
1.1	Trace and Determinants	5
1.2	The Special Case 2×2	6
2	Derivatives	7
2.1	Derivatives of a Determinant	7
2.2	Derivatives of an Inverse	8
2.3	Derivatives of Matrices, Vectors and Scalar Forms	9
2.4	Derivatives of Traces	11
2.5	Derivatives of Structured Matrices	12
3	Inverses	14
3.1	Exact Relations	14
3.2	Implication on Inverses	14
3.3	Approximations	15
3.4	Generalized Inverse	15
3.5	Pseudo Inverse	15
4	Complex Matrices	17
4.1	Complex Derivatives	17
5	Decompositions	20
5.1	Eigenvalues and Eigenvectors	20
5.2	Singular Value Decomposition	20
5.3	Triangular Decomposition	21
6	General Statistics and Probability	22
6.1	Moments of any distribution	22
6.2	Expectations	22
7	Gaussians	24
7.1	One Dimensional	24
7.2	Basics	25
7.3	Moments	27
7.4	Miscellaneous	29
7.5	One Dimensional Mixture of Gaussians	29
7.6	Mixture of Gaussians	30
8	Miscellaneous	31
8.1	Functions and Series	31
8.2	Indices, Entries and Vectors	32
8.3	Solutions to Systems of Equations	35
8.4	Block matrices	36
8.5	Matrix Norms	37
8.6	Positive Definite and Semi-definite Matrices	38

8.7	Integral Involving Dirac Delta Functions	39
8.8	Miscellaneous	40
A	Proofs and Details	41

Notation and Nomenclature

\mathbf{A}	Matrix
\mathbf{A}_{ij}	Matrix indexed for some purpose
\mathbf{A}_i	Matrix indexed for some purpose
\mathbf{A}^{ij}	Matrix indexed for some purpose
\mathbf{A}^n	Matrix indexed for some purpose or The n.th power of a square matrix
\mathbf{A}^{-1}	The inverse matrix of the matrix \mathbf{A}
\mathbf{A}^+	The pseudo inverse matrix of the matrix \mathbf{A}
$\mathbf{A}^{1/2}$	The square root of a matrix (if unique), not elementwise
$(\mathbf{A})_{ij}$	The (i, j) .th entry of the matrix \mathbf{A}
A_{ij}	The (i, j) .th entry of the matrix \mathbf{A}
\mathbf{a}	Vector
\mathbf{a}_i	Vector indexed for some purpose
a_i	The i.th element of the vector \mathbf{a}
a	Scalar
$\Re z$	Real part of a scalar
$\Re \mathbf{z}$	Real part of a vector
$\Re \mathbf{Z}$	Real part of a matrix
$\Im z$	Imaginary part of a scalar
$\Im \mathbf{z}$	Imaginary part of a vector
$\Im \mathbf{Z}$	Imaginary part of a matrix
$\det(\mathbf{A})$	Determinant of \mathbf{A}
$\ \mathbf{A}\ $	Matrix norm (subscript if any denotes what norm)
\mathbf{A}^T	Transposed matrix
\mathbf{A}^*	Complex conjugated matrix
\mathbf{A}^H	Transposed and complex conjugated matrix
$\mathbf{A} \circ \mathbf{B}$	Hadamard (elementwise) product
$\mathbf{A} \otimes \mathbf{B}$	Kronecker product
$\mathbf{0}$	The null matrix. Zero in all entries.
\mathbf{I}	The identity matrix
\mathbf{J}^{ij}	The single-entry matrix, 1 at (i, j) and zero elsewhere
Σ	A positive definite matrix
Λ	A diagonal matrix

1 Basics

$$\begin{aligned}(\mathbf{AB})^{-1} &= \mathbf{B}^{-1}\mathbf{A}^{-1} \\ (\mathbf{ABC}\dots)^{-1} &= \dots\mathbf{C}^{-1}\mathbf{B}^{-1}\mathbf{A}^{-1} \\ (\mathbf{A}^T)^{-1} &= (\mathbf{A}^{-1})^T \\ (\mathbf{A} + \mathbf{B})^T &= \mathbf{A}^T + \mathbf{B}^T \\ (\mathbf{AB})^T &= \mathbf{B}^T\mathbf{A}^T \\ (\mathbf{ABC}\dots)^T &= \dots\mathbf{C}^T\mathbf{B}^T\mathbf{A}^T \\ (\mathbf{A}^H)^{-1} &= (\mathbf{A}^{-1})^H \\ (\mathbf{A} + \mathbf{B})^H &= \mathbf{A}^H + \mathbf{B}^H \\ (\mathbf{AB})^H &= \mathbf{B}^H\mathbf{A}^H \\ (\mathbf{ABC}\dots)^H &= \dots\mathbf{C}^H\mathbf{B}^H\mathbf{A}^H\end{aligned}$$

1.1 Trace and Determinants

$$\begin{aligned}\text{Tr}(\mathbf{A}) &= \sum_i \mathbf{A}_{ii} = \sum_i \lambda_i, \quad \lambda_i = \text{eig}(\mathbf{A}) \\ \text{Tr}(\mathbf{A}) &= \text{Tr}(\mathbf{A}^T) \\ \text{Tr}(\mathbf{AB}) &= \text{Tr}(\mathbf{BA}) \\ \text{Tr}(\mathbf{A} + \mathbf{B}) &= \text{Tr}(\mathbf{A}) + \text{Tr}(\mathbf{B}) \\ \text{Tr}(\mathbf{ABC}) &= \text{Tr}(\mathbf{BCA}) = \text{Tr}(\mathbf{CAB}) \\ \det(\mathbf{A}) &= \prod_i \lambda_i \quad \lambda_i = \text{eig}(\mathbf{A}) \\ \det(\mathbf{AB}) &= \det(\mathbf{A})\det(\mathbf{B}), \quad \text{if } \mathbf{A} \text{ and } \mathbf{B} \text{ are invertible} \\ \det(\mathbf{A}^{-1}) &= \frac{1}{\det(\mathbf{A})}\end{aligned}$$

1.2 The Special Case 2x2

Consider the matrix \mathbf{A}

$$\mathbf{A} = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

Determinant and trace

$$\det(\mathbf{A}) = A_{11}A_{22} - A_{12}A_{21}$$

$$\text{Tr}(\mathbf{A}) = A_{11} + A_{22}$$

Eigenvalues

$$\lambda^2 - \lambda \cdot \text{Tr}(\mathbf{A}) + \det(\mathbf{A}) = 0$$

$$\lambda_1 = \frac{\text{Tr}(\mathbf{A}) + \sqrt{\text{Tr}(\mathbf{A})^2 - 4 \det(\mathbf{A})}}{2} \quad \lambda_2 = \frac{\text{Tr}(\mathbf{A}) - \sqrt{\text{Tr}(\mathbf{A})^2 - 4 \det(\mathbf{A})}}{2}$$

$$\lambda_1 + \lambda_2 = \text{Tr}(\mathbf{A}) \quad \lambda_1 \lambda_2 = \det(\mathbf{A})$$

Eigenvectors

$$\mathbf{v}_1 \propto \begin{bmatrix} A_{12} \\ \lambda_1 - A_{11} \end{bmatrix} \quad \mathbf{v}_2 \propto \begin{bmatrix} A_{12} \\ \lambda_2 - A_{11} \end{bmatrix}$$

Inverse

$$\mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \begin{bmatrix} A_{22} & -A_{12} \\ -A_{21} & A_{11} \end{bmatrix}$$

2 Derivatives

This section is covering differentiation of a number of expressions with respect to a matrix \mathbf{X} . Note that it is always assumed that \mathbf{X} has *no special structure*, i.e. that the elements of \mathbf{X} are independent (e.g. not symmetric, Toeplitz, positive definite). See section 2.5 for differentiation of structured matrices. The basic assumptions can be written in a formula as

$$\frac{\partial X_{kl}}{\partial X_{ij}} = \delta_{ik}\delta_{lj}$$

that is for e.g. vector forms,

$$\left[\frac{\partial \mathbf{x}}{\partial y} \right]_i = \frac{\partial x_i}{\partial y} \quad \left[\frac{\partial x}{\partial \mathbf{y}} \right]_i = \frac{\partial x}{\partial y_i} \quad \left[\frac{\partial \mathbf{x}}{\partial \mathbf{y}} \right]_{ij} = \frac{\partial x_i}{\partial y_j}$$

The following rules are general and very useful when deriving the differential of an expression ([10]):

$$\begin{aligned} \partial \mathbf{A} &= 0 && (\mathbf{A} \text{ is a constant}) && (1) \\ \partial(\alpha \mathbf{X}) &= \alpha \partial \mathbf{X} && && (2) \\ \partial(\mathbf{X} + \mathbf{Y}) &= \partial \mathbf{X} + \partial \mathbf{Y} && && (3) \\ \partial(\text{Tr}(\mathbf{X})) &= \text{Tr}(\partial \mathbf{X}) && && (4) \\ \partial(\mathbf{X}\mathbf{Y}) &= (\partial \mathbf{X})\mathbf{Y} + \mathbf{X}(\partial \mathbf{Y}) && && (5) \\ \partial(\mathbf{X} \circ \mathbf{Y}) &= (\partial \mathbf{X}) \circ \mathbf{Y} + \mathbf{X} \circ (\partial \mathbf{Y}) && && (6) \\ \partial(\mathbf{X} \otimes \mathbf{Y}) &= (\partial \mathbf{X}) \otimes \mathbf{Y} + \mathbf{X} \otimes (\partial \mathbf{Y}) && && (7) \\ \partial(\mathbf{X}^{-1}) &= -\mathbf{X}^{-1}(\partial \mathbf{X})\mathbf{X}^{-1} && && (8) \\ \partial(\det(\mathbf{X})) &= \det(\mathbf{X})\text{Tr}(\mathbf{X}^{-1}\partial \mathbf{X}) && && (9) \\ \partial(\ln(\det(\mathbf{X}))) &= \text{Tr}(\mathbf{X}^{-1}\partial \mathbf{X}) && && (10) \\ \partial \mathbf{X}^T &= (\partial \mathbf{X})^T && && (11) \\ \partial \mathbf{X}^H &= (\partial \mathbf{X})^H && && (12) \end{aligned}$$

2.1 Derivatives of a Determinant

2.1.1 General form

$$\frac{\partial \det(\mathbf{Y})}{\partial x} = \det(\mathbf{Y})\text{Tr} \left[\mathbf{Y}^{-1} \frac{\partial \mathbf{Y}}{\partial x} \right]$$

2.1.2 Linear forms

$$\begin{aligned} \frac{\partial \det(\mathbf{X})}{\partial \mathbf{X}} &= \det(\mathbf{X})(\mathbf{X}^{-1})^T \\ \frac{\partial \det(\mathbf{AXB})}{\partial \mathbf{X}} &= \det(\mathbf{AXB})(\mathbf{X}^{-1})^T = \det(\mathbf{AXB})(\mathbf{X}^T)^{-1} \end{aligned}$$

2.1.3 Square forms

If \mathbf{X} is square and invertible, then

$$\frac{\partial \det(\mathbf{X}^T \mathbf{A} \mathbf{X})}{\partial \mathbf{X}} = 2 \det(\mathbf{X}^T \mathbf{A} \mathbf{X}) \mathbf{X}^{-T}$$

If \mathbf{X} is not square but \mathbf{A} is symmetric, then

$$\frac{\partial \det(\mathbf{X}^T \mathbf{A} \mathbf{X})}{\partial \mathbf{X}} = 2 \det(\mathbf{X}^T \mathbf{A} \mathbf{X}) \mathbf{A} \mathbf{X} (\mathbf{X}^T \mathbf{A} \mathbf{X})^{-1}$$

If \mathbf{X} is not square and \mathbf{A} is not symmetric, then

$$\frac{\partial \det(\mathbf{X}^T \mathbf{A} \mathbf{X})}{\partial \mathbf{X}} = \det(\mathbf{X}^T \mathbf{A} \mathbf{X}) (\mathbf{A} \mathbf{X} (\mathbf{X}^T \mathbf{A} \mathbf{X})^{-1} + \mathbf{A}^T \mathbf{X} (\mathbf{X}^T \mathbf{A}^T \mathbf{X})^{-1}) \quad (13)$$

2.1.4 Other nonlinear forms

Some special cases are (See [8])

$$\begin{aligned} \frac{\partial \ln \det(\mathbf{X}^T \mathbf{X})}{\partial \mathbf{X}} &= 2(\mathbf{X}^+)^T \\ \frac{\partial \ln \det(\mathbf{X}^T \mathbf{X})}{\partial \mathbf{X}^+} &= -2\mathbf{X}^T \\ \frac{\partial \ln |\det(\mathbf{X})|}{\partial \mathbf{X}} &= (\mathbf{X}^{-1})^T = (\mathbf{X}^T)^{-1} \\ \frac{\partial \det(\mathbf{X}^k)}{\partial \mathbf{X}} &= k \det(\mathbf{X}^k) \mathbf{X}^{-T} \end{aligned}$$

See [7].

2.2 Derivatives of an Inverse

From [15] we have the basic identity

$$\frac{\partial \mathbf{Y}^{-1}}{\partial x} = -\mathbf{Y}^{-1} \frac{\partial \mathbf{Y}}{\partial x} \mathbf{Y}^{-1}$$

from which it follows

$$\begin{aligned} \frac{\partial (\mathbf{X}^{-1})_{kl}}{\partial X_{ij}} &= -(\mathbf{X}^{-1})_{ki} (\mathbf{X}^{-1})_{jl} \\ \frac{\partial \mathbf{a}^T \mathbf{X}^{-1} \mathbf{b}}{\partial \mathbf{X}} &= -\mathbf{X}^{-T} \mathbf{a} \mathbf{b}^T \mathbf{X}^{-T} \\ \frac{\partial \det(\mathbf{X}^{-1})}{\partial \mathbf{X}} &= -\det(\mathbf{X}^{-1}) (\mathbf{X}^{-1})^T \\ \frac{\partial \text{Tr}(\mathbf{A} \mathbf{X}^{-1} \mathbf{B})}{\partial \mathbf{X}} &= -(\mathbf{X}^{-1} \mathbf{B} \mathbf{A} \mathbf{X}^{-1})^T \end{aligned}$$

2.3 Derivatives of Matrices, Vectors and Scalar Forms

2.3.1 First Order

$$\begin{aligned} \frac{\partial \mathbf{x}^T \mathbf{b}}{\partial \mathbf{x}} &= \frac{\partial \mathbf{b}^T \mathbf{x}}{\partial \mathbf{x}} = \mathbf{b} \\ \frac{\partial \mathbf{a}^T \mathbf{X} \mathbf{b}}{\partial \mathbf{X}} &= \mathbf{a} \mathbf{b}^T \\ \frac{\partial \mathbf{a}^T \mathbf{X}^T \mathbf{b}}{\partial \mathbf{X}} &= \mathbf{b} \mathbf{a}^T \\ \frac{\partial \mathbf{a}^T \mathbf{X} \mathbf{a}}{\partial \mathbf{X}} &= \frac{\partial \mathbf{a}^T \mathbf{X}^T \mathbf{a}}{\partial \mathbf{X}} = \mathbf{a} \mathbf{a}^T \\ \frac{\partial \mathbf{X}}{\partial X_{ij}} &= \mathbf{J}^{ij} \\ \frac{\partial (\mathbf{X} \mathbf{A})_{ij}}{\partial X_{mn}} &= \delta_{im} (\mathbf{A})_{nj} = (\mathbf{J}^{mn} \mathbf{A})_{ij} \\ \frac{\partial (\mathbf{X}^T \mathbf{A})_{ij}}{\partial X_{mn}} &= \delta_{in} (\mathbf{A})_{mj} = (\mathbf{J}^{nm} \mathbf{A})_{ij} \end{aligned}$$

2.3.2 Second Order

$$\begin{aligned} \frac{\partial}{\partial X_{ij}} \sum_{klmn} X_{kl} X_{mn} &= 2 \sum_{kl} X_{kl} \\ \frac{\partial \mathbf{b}^T \mathbf{X}^T \mathbf{X} \mathbf{c}}{\partial \mathbf{X}} &= \mathbf{X} (\mathbf{b} \mathbf{c}^T + \mathbf{c} \mathbf{b}^T) \\ \frac{\partial (\mathbf{B} \mathbf{x} + \mathbf{b})^T \mathbf{C} (\mathbf{D} \mathbf{x} + \mathbf{d})}{\partial \mathbf{x}} &= \mathbf{B}^T \mathbf{C} (\mathbf{D} \mathbf{x} + \mathbf{d}) + \mathbf{D}^T \mathbf{C}^T (\mathbf{B} \mathbf{x} + \mathbf{b}) \\ \frac{\partial (\mathbf{X}^T \mathbf{B} \mathbf{X})_{kl}}{\partial X_{ij}} &= \delta_{lj} (\mathbf{X}^T \mathbf{B})_{ki} + \delta_{kj} (\mathbf{B} \mathbf{X})_{il} \\ \frac{\partial (\mathbf{X}^T \mathbf{B} \mathbf{X})}{\partial X_{ij}} &= \mathbf{X}^T \mathbf{B} \mathbf{J}^{ij} + \mathbf{J}^{ji} \mathbf{B} \mathbf{X} \quad (\mathbf{J}^{ij})_{kl} = \delta_{ik} \delta_{jl} \end{aligned}$$

See Sec 8.2 for useful properties of the Single-entry matrix \mathbf{J}^{ij}

$$\begin{aligned} \frac{\partial \mathbf{x}^T \mathbf{B} \mathbf{x}}{\partial \mathbf{x}} &= (\mathbf{B} + \mathbf{B}^T) \mathbf{x} \\ \frac{\partial \mathbf{b}^T \mathbf{X}^T \mathbf{D} \mathbf{X} \mathbf{c}}{\partial \mathbf{X}} &= \mathbf{D}^T \mathbf{X} \mathbf{b} \mathbf{c}^T + \mathbf{D} \mathbf{X} \mathbf{c} \mathbf{b}^T \\ \frac{\partial}{\partial \mathbf{X}} (\mathbf{X} \mathbf{b} + \mathbf{c})^T \mathbf{D} (\mathbf{X} \mathbf{b} + \mathbf{c}) &= (\mathbf{D} + \mathbf{D}^T) (\mathbf{X} \mathbf{b} + \mathbf{c}) \mathbf{b}^T \end{aligned}$$

Assume \mathbf{W} is symmetric, then

$$\begin{aligned}\frac{\partial}{\partial \mathbf{s}}(\mathbf{x} - \mathbf{A}\mathbf{s})^T \mathbf{W}(\mathbf{x} - \mathbf{A}\mathbf{s}) &= -2\mathbf{A}^T \mathbf{W}(\mathbf{x} - \mathbf{A}\mathbf{s}) \\ \frac{\partial}{\partial \mathbf{x}}(\mathbf{x} - \mathbf{A}\mathbf{s})^T \mathbf{W}(\mathbf{x} - \mathbf{A}\mathbf{s}) &= -2\mathbf{W}(\mathbf{x} - \mathbf{A}\mathbf{s}) \\ \frac{\partial}{\partial \mathbf{A}}(\mathbf{x} - \mathbf{A}\mathbf{s})^T \mathbf{W}(\mathbf{x} - \mathbf{A}\mathbf{s}) &= -2\mathbf{W}(\mathbf{x} - \mathbf{A}\mathbf{s})\mathbf{s}^T\end{aligned}$$

2.3.3 Higher order and non-linear

$$\frac{\partial}{\partial \mathbf{X}} \mathbf{a}^T \mathbf{X}^n \mathbf{b} = \sum_{r=0}^{n-1} (\mathbf{X}^r)^T \mathbf{a} \mathbf{b}^T (\mathbf{X}^{n-1-r})^T \quad (14)$$

$$\begin{aligned}\frac{\partial}{\partial \mathbf{X}} \mathbf{a}^T (\mathbf{X}^n)^T \mathbf{X}^n \mathbf{b} &= \sum_{r=0}^{n-1} \left[\mathbf{X}^{n-1-r} \mathbf{a} \mathbf{b}^T (\mathbf{X}^n)^T \mathbf{X}^r \right. \\ &\quad \left. + (\mathbf{X}^r)^T \mathbf{X}^n \mathbf{a} \mathbf{b}^T (\mathbf{X}^{n-1-r})^T \right] \quad (15)\end{aligned}$$

See A.0.1 for a proof.

Assume \mathbf{s} and \mathbf{r} are functions of \mathbf{x} , i.e. $\mathbf{s} = \mathbf{s}(\mathbf{x})$, $\mathbf{r} = \mathbf{r}(\mathbf{x})$, and that \mathbf{A} is a constant, then

$$\begin{aligned}\frac{\partial}{\partial \mathbf{x}} \mathbf{s}^T \mathbf{A} \mathbf{s} &= \left[\frac{\partial \mathbf{s}}{\partial \mathbf{x}} \right]^T (\mathbf{A} + \mathbf{A}^T) \mathbf{s} \\ \frac{\partial}{\partial \mathbf{x}} \mathbf{s}^T \mathbf{A} \mathbf{r} &= \left[\frac{\partial \mathbf{s}}{\partial \mathbf{x}} \right]^T \mathbf{A} \mathbf{s} + \left[\frac{\partial \mathbf{r}}{\partial \mathbf{x}} \right]^T \mathbf{A}^T \mathbf{r}\end{aligned}$$

2.3.4 Gradient and Hessian

Using the above we have for the gradient and the hessian

$$\begin{aligned}f &= \mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} \\ \nabla_{\mathbf{x}} f &= \frac{\partial f}{\partial \mathbf{x}} = (\mathbf{A} + \mathbf{A}^T) \mathbf{x} + \mathbf{b} \\ \frac{\partial^2 f}{\partial \mathbf{x} \partial \mathbf{x}^T} &= \mathbf{A} + \mathbf{A}^T\end{aligned}$$

2.4 Derivatives of Traces

2.4.1 First Order

$$\begin{aligned}
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{X}) &= \mathbf{I} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{XB}) &= \mathbf{B}^T \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{BXC}) &= \mathbf{B}^T \mathbf{C}^T \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{BX}^T \mathbf{C}) &= \mathbf{CB} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{X}^T \mathbf{C}) &= \mathbf{C} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{BX}^T) &= \mathbf{B}
 \end{aligned} \tag{16}$$

2.4.2 Second Order

$$\begin{aligned}
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{X}^2) &= 2\mathbf{X} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{X}^2 \mathbf{B}) &= (\mathbf{XB} + \mathbf{BX})^T \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{X}^T \mathbf{BX}) &= \mathbf{BX} + \mathbf{B}^T \mathbf{X} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{XBX}^T) &= \mathbf{XB}^T + \mathbf{XB} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{X}^T \mathbf{X}) &= 2\mathbf{X} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{BXX}^T) &= (\mathbf{B} + \mathbf{B}^T) \mathbf{X} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{B}^T \mathbf{X}^T \mathbf{CXB}) &= \mathbf{C}^T \mathbf{XBB}^T + \mathbf{CXBB}^T \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}[\mathbf{X}^T \mathbf{BXC}] &= \mathbf{BXC} + \mathbf{B}^T \mathbf{XC}^T \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{AXBX}^T \mathbf{C}) &= \mathbf{A}^T \mathbf{C}^T \mathbf{XB}^T + \mathbf{CAXB} \\
 \frac{\partial}{\partial \mathbf{X}} \text{Tr}[(\mathbf{AXb} + \mathbf{c})(\mathbf{AXb} + \mathbf{c})^T] &= 2\mathbf{A}^T (\mathbf{AXb} + \mathbf{c}) \mathbf{b}^T
 \end{aligned}$$

See [7].

2.4.3 Higher Order

$$\frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{X}^k) = k(\mathbf{X}^{k-1})^T$$

$$\frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{A}\mathbf{X}^k) = \sum_{r=0}^{k-1} (\mathbf{X}^r \mathbf{A} \mathbf{X}^{k-r-1})^T$$

$$\begin{aligned} \frac{\partial}{\partial \mathbf{X}} \text{Tr}[\mathbf{B}^T \mathbf{X}^T \mathbf{C} \mathbf{X} \mathbf{X}^T \mathbf{C} \mathbf{X} \mathbf{B}] &= \mathbf{C} \mathbf{X} \mathbf{X}^T \mathbf{C} \mathbf{X} \mathbf{B} \mathbf{B}^T + \mathbf{C}^T \mathbf{X} \mathbf{B} \mathbf{B}^T \mathbf{X}^T \mathbf{C}^T \mathbf{X} \\ &+ \mathbf{C} \mathbf{X} \mathbf{B} \mathbf{B}^T \mathbf{X}^T \mathbf{C} \mathbf{X} + \mathbf{C}^T \mathbf{X} \mathbf{X}^T \mathbf{C}^T \mathbf{X} \mathbf{B} \mathbf{B}^T \end{aligned}$$

2.4.4 Other

$$\frac{\partial}{\partial \mathbf{X}} \text{Tr}(\mathbf{A}\mathbf{X}^{-1}\mathbf{B}) = -(\mathbf{X}^{-1}\mathbf{B}\mathbf{A}\mathbf{X}^{-1})^T = -\mathbf{X}^{-T} \mathbf{A}^T \mathbf{B}^T \mathbf{X}^{-T}$$

Assume \mathbf{B} and \mathbf{C} to be symmetric, then

$$\frac{\partial}{\partial \mathbf{X}} \text{Tr}[(\mathbf{X}^T \mathbf{C} \mathbf{X})^{-1} \mathbf{A}] = -(\mathbf{C} \mathbf{X} (\mathbf{X}^T \mathbf{C} \mathbf{X})^{-1}) (\mathbf{A} + \mathbf{A}^T) (\mathbf{X}^T \mathbf{C} \mathbf{X})^{-1}$$

$$\begin{aligned} \frac{\partial}{\partial \mathbf{X}} \text{Tr}[(\mathbf{X}^T \mathbf{C} \mathbf{X})^{-1} (\mathbf{X}^T \mathbf{B} \mathbf{X})] &= -2\mathbf{C} \mathbf{X} (\mathbf{X}^T \mathbf{C} \mathbf{X})^{-1} \mathbf{X}^T \mathbf{B} \mathbf{X} (\mathbf{X}^T \mathbf{C} \mathbf{X})^{-1} \\ &+ 2\mathbf{B} \mathbf{X} (\mathbf{X}^T \mathbf{C} \mathbf{X})^{-1} \end{aligned}$$

See [7].

2.5 Derivatives of Structured Matrices

Assume that the matrix \mathbf{A} has some structure, i.e. is symmetric, toeplitz, etc. In that case the derivatives of the previous section does not apply in general. In stead, consider the following general rule for differentiating a scalar function $f(\mathbf{A})$

$$\frac{df}{dA_{ij}} = \sum_{kl} \frac{\partial f}{\partial A_{kl}} \frac{\partial A_{kl}}{\partial A_{ij}} = \text{Tr} \left[\left[\frac{\partial f}{\partial \mathbf{A}} \right]^T \frac{\partial \mathbf{A}}{\partial A_{ij}} \right]$$

The matrix differentiated with respect to itself is in this document referred to as the *structure matrix* of \mathbf{A} and is defined simply by

$$\frac{\partial \mathbf{A}}{\partial A_{ij}} = \mathbf{S}^{ij}$$

If \mathbf{A} has no special structure we have simply $\mathbf{S}^{ij} = \mathbf{J}^{ij}$, that is, the structure matrix is simply the singleentry matrix. Many structures have a representation in singleentry matrices, see Sec. 8.2.7 for more examples of structure matrices.

2.5.1 Symmetric

If \mathbf{A} is symmetric, then $\mathbf{S}^{ij} = \mathbf{J}^{ij} + \mathbf{J}^{ji} - \mathbf{J}^{ij}\mathbf{J}^{ij}$ and therefore

$$\frac{df}{d\mathbf{A}} = \left[\frac{\partial f}{\partial \mathbf{A}} \right] + \left[\frac{\partial f}{\partial \mathbf{A}} \right]^T - \text{diag} \left[\frac{\partial f}{\partial \mathbf{A}} \right]$$

That is, e.g., ([5], [16]):

$$\frac{\partial \text{Tr}(\mathbf{A}\mathbf{X})}{\partial \mathbf{X}} = \mathbf{A} + \mathbf{A}^T - (\mathbf{A} \circ \mathbf{I}), \text{ see (20)} \quad (17)$$

$$\frac{\partial \det(\mathbf{X})}{\partial \mathbf{X}} = 2\mathbf{X} - (\mathbf{X} \circ \mathbf{I}) \quad (18)$$

$$\frac{\partial \ln \det(\mathbf{X})}{\partial \mathbf{X}} = 2\mathbf{X}^{-1} - (\mathbf{X}^{-1} \circ \mathbf{I}) \quad (19)$$

2.5.2 Diagonal

If \mathbf{X} is diagonal, then ([10]):

$$\frac{\partial \text{Tr}(\mathbf{A}\mathbf{X})}{\partial \mathbf{X}} = \mathbf{A} \circ \mathbf{I} \quad (20)$$

3 Inverses

3.1 Exact Relations

3.1.1 The Woodbury identity

$$(\mathbf{A} + \mathbf{C}\mathbf{B}\mathbf{C}^T)^{-1} = \mathbf{A}^{-1} - \mathbf{A}^{-1}\mathbf{C}(\mathbf{B}^{-1} + \mathbf{C}^T\mathbf{A}^{-1}\mathbf{C})^{-1}\mathbf{C}^T\mathbf{A}^{-1}$$

If \mathbf{P}, \mathbf{R} are positive definite, then (see [17])

$$(\mathbf{P}^{-1} + \mathbf{B}^T\mathbf{R}^{-1}\mathbf{B})^{-1}\mathbf{B}^T\mathbf{R}^{-1} = \mathbf{P}\mathbf{B}^T(\mathbf{B}\mathbf{P}\mathbf{B}^T + \mathbf{R})^{-1}$$

3.1.2 The Kailath Variant

$$(\mathbf{A} + \mathbf{B}\mathbf{C})^{-1} = \mathbf{A}^{-1} - \mathbf{A}^{-1}\mathbf{B}(\mathbf{I} + \mathbf{C}\mathbf{A}^{-1}\mathbf{B})^{-1}\mathbf{C}\mathbf{A}^{-1}$$

See [4] page 153.

3.1.3 The Searle Set of Identities

The following set of identities, can be found in [13], page 151,

$$(\mathbf{I} + \mathbf{A}^{-1})^{-1} = \mathbf{A}(\mathbf{A} + \mathbf{I})^{-1}$$

$$(\mathbf{A} + \mathbf{B}\mathbf{B}^T)^{-1}\mathbf{B} = \mathbf{A}^{-1}\mathbf{B}(\mathbf{I} + \mathbf{B}^T\mathbf{A}^{-1}\mathbf{B})^{-1}$$

$$(\mathbf{A}^{-1} + \mathbf{B}^{-1})^{-1} = \mathbf{A}(\mathbf{A} + \mathbf{B})^{-1}\mathbf{B} = \mathbf{B}(\mathbf{A} + \mathbf{B})^{-1}\mathbf{A}$$

$$\mathbf{A} - \mathbf{A}(\mathbf{A} + \mathbf{B})^{-1}\mathbf{A} = \mathbf{B} - \mathbf{B}(\mathbf{A} + \mathbf{B})^{-1}\mathbf{B}$$

$$\mathbf{A}^{-1} + \mathbf{B}^{-1} = \mathbf{A}^{-1}(\mathbf{A} + \mathbf{B})\mathbf{B}^{-1}$$

$$(\mathbf{I} + \mathbf{A}\mathbf{B})^{-1} = \mathbf{I} - \mathbf{A}(\mathbf{I} + \mathbf{B}\mathbf{A})^{-1}\mathbf{B}$$

$$(\mathbf{I} + \mathbf{A}\mathbf{B})^{-1}\mathbf{A} = \mathbf{A}(\mathbf{I} + \mathbf{A}\mathbf{B})^{-1}$$

3.2 Implication on Inverses

$$(\mathbf{A} + \mathbf{B})^{-1} = \mathbf{A}^{-1} + \mathbf{B}^{-1} \quad \Rightarrow \quad \mathbf{A}\mathbf{B}^{-1}\mathbf{A} = \mathbf{B}\mathbf{A}^{-1}\mathbf{B}$$

See [13].

3.2.1 A PosDef identity

Assume \mathbf{P}, \mathbf{R} to be positive definite and invertible, then

$$(\mathbf{P}^{-1} + \mathbf{B}^T\mathbf{R}^{-1}\mathbf{B})^{-1}\mathbf{B}^T\mathbf{R}^{-1} = \mathbf{P}\mathbf{B}^T(\mathbf{B}\mathbf{P}\mathbf{B}^T + \mathbf{R})^{-1}$$

See [?].

3.3 Approximations

$$(\mathbf{I} + \mathbf{A})^{-1} = \mathbf{I} - \mathbf{A} + \mathbf{A}^2 - \mathbf{A}^3 + \dots$$

$$\mathbf{A} - \mathbf{A}(\mathbf{I} + \mathbf{A})^{-1}\mathbf{A} \cong \mathbf{I} - \mathbf{A}^{-1} \quad \text{if } \mathbf{A} \text{ large and symmetric}$$

If σ^2 is small then

$$(\mathbf{Q} + \sigma^2\mathbf{M})^{-1} \cong \mathbf{Q}^{-1} - \sigma^2\mathbf{Q}^{-1}\mathbf{M}\mathbf{Q}^{-1}$$

3.4 Generalized Inverse

3.4.1 Definition

A generalized inverse matrix of the matrix \mathbf{A} is any matrix \mathbf{A}^- such that

$$\mathbf{A}\mathbf{A}^-\mathbf{A} = \mathbf{A}$$

The matrix \mathbf{A}^- is not unique.

3.5 Pseudo Inverse

3.5.1 Definition

The pseudo inverse (or Moore-Penrose inverse) of a matrix \mathbf{A} is the matrix \mathbf{A}^+ that fulfils

$$\begin{array}{ll} \text{I} & \mathbf{A}\mathbf{A}^+\mathbf{A} = \mathbf{A} \\ \text{II} & \mathbf{A}^+\mathbf{A}\mathbf{A}^+ = \mathbf{A}^+ \\ \text{III} & \mathbf{A}\mathbf{A}^+ \text{ symmetric} \\ \text{IV} & \mathbf{A}^+\mathbf{A} \text{ symmetric} \end{array}$$

The matrix \mathbf{A}^+ is unique and does always exist.

3.5.2 Properties

Assume \mathbf{A}^+ to be the pseudo-inverse of \mathbf{A} , then (See [3])

$$\begin{array}{ll} (\mathbf{A}^+)^+ & = \mathbf{A} \\ (\mathbf{A}^T)^+ & = (\mathbf{A}^+)^T \\ (c\mathbf{A})^+ & = (1/c)\mathbf{A}^+ \\ (\mathbf{A}^T\mathbf{A})^+ & = \mathbf{A}^+(\mathbf{A}^T)^+ \\ (\mathbf{A}\mathbf{A}^T)^+ & = (\mathbf{A}^T)^+\mathbf{A}^+ \end{array}$$

Assume \mathbf{A} to have full rank, then

$$\begin{array}{ll} (\mathbf{A}\mathbf{A}^+)(\mathbf{A}\mathbf{A}^+) & = \mathbf{A}\mathbf{A}^+ \\ (\mathbf{A}^+\mathbf{A})(\mathbf{A}^+\mathbf{A}) & = \mathbf{A}^+\mathbf{A} \\ \text{Tr}(\mathbf{A}\mathbf{A}^+) & = \text{rank}(\mathbf{A}\mathbf{A}^+) \quad (\text{See [14]}) \\ \text{Tr}(\mathbf{A}^+\mathbf{A}) & = \text{rank}(\mathbf{A}^+\mathbf{A}) \quad (\text{See [14]}) \end{array}$$

3.5.3 Construction

Assume that \mathbf{A} has full rank, then

$$\begin{array}{llll} \mathbf{A} \ n \times n & \text{Square} & \text{rank}(\mathbf{A}) = n & \Rightarrow \mathbf{A}^+ = \mathbf{A}^{-1} \\ \mathbf{A} \ n \times m & \text{Broad} & \text{rank}(\mathbf{A}) = n & \Rightarrow \mathbf{A}^+ = \mathbf{A}^T(\mathbf{A}\mathbf{A}^T)^{-1} \\ \mathbf{A} \ n \times m & \text{Tall} & \text{rank}(\mathbf{A}) = m & \Rightarrow \mathbf{A}^+ = (\mathbf{A}^T\mathbf{A})^{-1}\mathbf{A}^T \end{array}$$

Assume \mathbf{A} does not have full rank, i.e. \mathbf{A} is $n \times m$ and $\text{rank}(\mathbf{A}) = r < \min(n, m)$. The pseudo inverse \mathbf{A}^+ can be constructed from the singular value decomposition $\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$, by

$$\mathbf{A}^+ = \mathbf{V}\mathbf{D}^+\mathbf{U}^T$$

A different way is this: There does always exist two matrices \mathbf{C} $n \times r$ and \mathbf{D} $r \times m$ of rank r , such that $\mathbf{A} = \mathbf{C}\mathbf{D}$. Using these matrices it holds that

$$\mathbf{A}^+ = \mathbf{D}^T(\mathbf{D}\mathbf{D}^T)^{-1}(\mathbf{C}^T\mathbf{C})^{-1}\mathbf{C}^T$$

See [3].

4 Complex Matrices

4.1 Complex Derivatives

In order to differentiate an expression $f(z)$ with respect to a complex z , the Cauchy-Riemann equations have to be satisfied ([7]):

$$\frac{df(z)}{dz} = \frac{\partial \Re(f(z))}{\partial \Re z} + i \frac{\partial \Im(f(z))}{\partial \Re z} \quad (21)$$

and

$$\frac{df(z)}{dz} = -i \frac{\partial \Re(f(z))}{\partial \Im z} + \frac{\partial \Im(f(z))}{\partial \Im z} \quad (22)$$

or in a more compact form:

$$\frac{\partial f(z)}{\partial \Im z} = i \frac{\partial f(z)}{\partial \Re z}. \quad (23)$$

A complex function that satisfies the Cauchy-Riemann equations for points in a region R is said to be *analytic* in this region R . In general, expressions involving complex conjugate or conjugate transpose do not satisfy the Cauchy-Riemann equations. In order to avoid this problem, a more generalized definition of complex derivative is used ([12], [6]):

- Generalized Complex Derivative:

$$\frac{df(z)}{dz} = \frac{1}{2} \left(\frac{\partial f(z)}{\partial \Re z} - i \frac{\partial f(z)}{\partial \Im z} \right) \quad (24)$$

- Conjugate Complex Derivative

$$\frac{df(z)}{dz^*} = \frac{1}{2} \left(\frac{\partial f(z)}{\partial \Re z} + i \frac{\partial f(z)}{\partial \Im z} \right) \quad (25)$$

The Generalized Complex Derivative equals the normal derivative, when f is an analytic function. For a non-analytic function such as $f(z) = z^*$, the derivative equals zero. The Conjugate Complex Derivative equals zero, when f is an analytic function. The Conjugate Complex Derivative has e.g. been used by [11] when deriving a complex gradient.

Notice:

$$\frac{df(z)}{dz} \neq \frac{\partial f(z)}{\partial \Re z} + i \frac{\partial f(z)}{\partial \Im z} \quad (26)$$

- Complex Gradient Vector: If f is a real function of a complex vector \mathbf{z} , then the complex gradient vector is given by ([9, p. 798])

$$\begin{aligned} \nabla f(\mathbf{z}) &= 2 \frac{df(\mathbf{z})}{d\mathbf{z}^*} \\ &= \frac{\partial f(\mathbf{z})}{\partial \Re \mathbf{z}} + i \frac{\partial f(\mathbf{z})}{\partial \Im \mathbf{z}} \end{aligned} \quad (27)$$

- **Complex Gradient Matrix:** If f is a real function of a complex matrix \mathbf{Z} , then the complex gradient matrix is given by ([2])

$$\begin{aligned}\nabla f(\mathbf{Z}) &= 2 \frac{df(\mathbf{Z})}{d\mathbf{Z}^*} \\ &= \frac{\partial f(\mathbf{Z})}{\partial \Re \mathbf{Z}} + i \frac{\partial f(\mathbf{Z})}{\partial \Im \mathbf{Z}}\end{aligned}\quad (28)$$

These expressions can be used for gradient descent algorithms.

4.1.1 The Chain Rule for complex numbers

The chain rule is a little more complicated when the function of a complex $u = f(x)$ is non-analytic. For a non-analytic function, the following chain rule can be applied ([?])

$$\begin{aligned}\frac{\partial g(u)}{\partial x} &= \frac{\partial g}{\partial u} \frac{\partial u}{\partial x} + \frac{\partial g}{\partial u^*} \frac{\partial u^*}{\partial x} \\ &= \frac{\partial g}{\partial u} \frac{\partial u}{\partial x} + \left(\frac{\partial g}{\partial u} \right)^* \frac{\partial u^*}{\partial x}\end{aligned}\quad (29)$$

Notice, if the function is analytic, the second term reduces to zero, and the function is reduced to the normal well-known chain rule. For the matrix derivative of a scalar function $g(\mathbf{U})$, the chain rule can be written the following way:

$$\frac{\partial g(\mathbf{U})}{\partial \mathbf{X}} = \frac{\text{Tr}\left(\left(\frac{\partial g(\mathbf{U})}{\partial \mathbf{U}}\right)^T \partial \mathbf{U}\right)}{\partial \mathbf{X}} + \frac{\text{Tr}\left(\left(\frac{\partial g(\mathbf{U})}{\partial \mathbf{U}^*}\right)^T \partial \mathbf{U}^*\right)}{\partial \mathbf{X}}. \quad (30)$$

4.1.2 Complex Derivatives of Traces

If the derivatives involve complex numbers, the conjugate transpose is often involved. The most useful way to show complex derivative is to show the derivative with respect to the real and the imaginary part separately. An easy example is:

$$\frac{\partial \text{Tr}(\mathbf{X}^*)}{\partial \Re \mathbf{X}} = \frac{\partial \text{Tr}(\mathbf{X}^H)}{\partial \Re \mathbf{X}} = \mathbf{I} \quad (31)$$

$$i \frac{\partial \text{Tr}(\mathbf{X}^*)}{\partial \Im \mathbf{X}} = i \frac{\partial \text{Tr}(\mathbf{X}^H)}{\partial \Im \mathbf{X}} = \mathbf{I} \quad (32)$$

Since the two results have the same sign, the conjugate complex derivative (25) should be used.

$$\frac{\partial \text{Tr}(\mathbf{X})}{\partial \Re \mathbf{X}} = \frac{\partial \text{Tr}(\mathbf{X}^T)}{\partial \Re \mathbf{X}} = \mathbf{I} \quad (33)$$

$$i \frac{\partial \text{Tr}(\mathbf{X})}{\partial \Im \mathbf{X}} = i \frac{\partial \text{Tr}(\mathbf{X}^T)}{\partial \Im \mathbf{X}} = -\mathbf{I} \quad (34)$$

Here, the two results have different signs, the generalized complex derivative (24) should be used. Hereby, it can be seen that (??) holds even if \mathbf{X} is a

complex number.

$$\frac{\partial \text{Tr}(\mathbf{A}\mathbf{X}^H)}{\partial \Re \mathbf{X}} = \mathbf{A} \quad (35)$$

$$i \frac{\partial \text{Tr}(\mathbf{A}\mathbf{X}^H)}{\partial \Im \mathbf{X}} = \mathbf{A} \quad (36)$$

$$\frac{\partial \text{Tr}(\mathbf{A}\mathbf{X}^*)}{\partial \Re \mathbf{X}} = \mathbf{A}^T \quad (37)$$

$$i \frac{\partial \text{Tr}(\mathbf{A}\mathbf{X}^*)}{\partial \Im \mathbf{X}} = \mathbf{A}^T \quad (38)$$

$$\frac{\partial \text{Tr}(\mathbf{X}\mathbf{X}^H)}{\partial \Re \mathbf{X}} = \frac{\partial \text{Tr}(\mathbf{X}^H \mathbf{X})}{\partial \Re \mathbf{X}} = 2\Re \mathbf{X} \quad (39)$$

$$i \frac{\partial \text{Tr}(\mathbf{X}\mathbf{X}^H)}{\partial \Im \mathbf{X}} = i \frac{\partial \text{Tr}(\mathbf{X}^H \mathbf{X})}{\partial \Im \mathbf{X}} = i2\Im \mathbf{X} \quad (40)$$

By inserting (39) and (40) in (24) and (25), it can be seen that

$$\frac{\partial \text{Tr}(\mathbf{X}\mathbf{X}^H)}{\partial \mathbf{X}} = \mathbf{X}^* \quad (41)$$

$$\frac{\partial \text{Tr}(\mathbf{X}\mathbf{X}^H)}{\partial \mathbf{X}^*} = \mathbf{X} \quad (42)$$

Since the function $\text{Tr}(\mathbf{X}\mathbf{X}^H)$ is a real function of the complex matrix \mathbf{X} , the complex gradient matrix (28) is given by

$$\nabla \text{Tr}(\mathbf{X}\mathbf{X}^H) = 2 \frac{\partial \text{Tr}(\mathbf{X}\mathbf{X}^H)}{\partial \mathbf{X}^*} = 2\mathbf{X} \quad (43)$$

4.1.3 Complex Derivative Involving Determinants

Here, a calculation example is provided. The objective is to find the derivative of $\det(\mathbf{X}^H \mathbf{A}\mathbf{X})$ with respect to $\mathbf{X} \in \mathbb{C}^{m \times n}$. The derivative is found with respect to the real part and the imaginary part of \mathbf{X} , by use of (9) and (5), $\det(\mathbf{X}^H \mathbf{A}\mathbf{X})$ can be calculated as (see Sec. A.0.2 for details)

$$\begin{aligned} \frac{\partial \det(\mathbf{X}^H \mathbf{A}\mathbf{X})}{\partial \mathbf{X}} &= \frac{1}{2} \left(\frac{\partial \det(\mathbf{X}^H \mathbf{A}\mathbf{X})}{\partial \Re \mathbf{X}} - i \frac{\partial \det(\mathbf{X}^H \mathbf{A}\mathbf{X})}{\partial \Im \mathbf{X}} \right) \\ &= \det(\mathbf{X}^H \mathbf{A}\mathbf{X}) ((\mathbf{X}^H \mathbf{A}\mathbf{X})^{-1} \mathbf{X}^H \mathbf{A})^T \end{aligned} \quad (44)$$

and the complex conjugate derivative yields

$$\begin{aligned} \frac{\partial \det(\mathbf{X}^H \mathbf{A}\mathbf{X})}{\partial \mathbf{X}^*} &= \frac{1}{2} \left(\frac{\partial \det(\mathbf{X}^H \mathbf{A}\mathbf{X})}{\partial \Re \mathbf{X}} + i \frac{\partial \det(\mathbf{X}^H \mathbf{A}\mathbf{X})}{\partial \Im \mathbf{X}} \right) \\ &= \det(\mathbf{X}^H \mathbf{A}\mathbf{X}) \mathbf{A}\mathbf{X} (\mathbf{X}^H \mathbf{A}\mathbf{X})^{-1} \end{aligned} \quad (45)$$

5 Decompositions

5.1 Eigenvalues and Eigenvectors

5.1.1 Definition

The eigenvectors \mathbf{v} and eigenvalues λ are the ones satisfying

$$\begin{aligned}\mathbf{A}\mathbf{v}_i &= \lambda_i\mathbf{v}_i \\ \mathbf{A}\mathbf{V} &= \mathbf{V}\mathbf{D}, \quad (\mathbf{D})_{ij} = \delta_{ij}\lambda_i\end{aligned}$$

where the columns of \mathbf{V} are the vectors \mathbf{v}_i

5.1.2 General Properties

$$\begin{aligned}\text{eig}(\mathbf{A}\mathbf{B}) &= \text{eig}(\mathbf{B}\mathbf{A}) \\ \mathbf{A} \text{ is } n \times m &\Rightarrow \text{At most } \min(n, m) \text{ distinct } \lambda_i \\ \text{rank}(\mathbf{A}) = r &\Rightarrow \text{At most } r \text{ non-zero } \lambda_i\end{aligned}$$

5.1.3 Symmetric

Assume \mathbf{A} is symmetric, then

$$\begin{aligned}\mathbf{V}\mathbf{V}^T &= \mathbf{I} \quad (\text{i.e. } \mathbf{V} \text{ is orthogonal}) \\ \lambda_i &\in \mathbb{R} \quad (\text{i.e. } \lambda_i \text{ is real}) \\ \text{Tr}(\mathbf{A}^p) &= \sum_i \lambda_i^p \\ \text{eig}(\mathbf{I} + c\mathbf{A}) &= 1 + c\lambda_i \\ \text{eig}(\mathbf{A}^{-1}) &= \lambda_i^{-1}\end{aligned}$$

5.2 Singular Value Decomposition

Any $n \times m$ matrix \mathbf{A} can be written as

$$\mathbf{A} = \mathbf{U}\mathbf{D}\mathbf{V}^T$$

where

$$\begin{aligned}\mathbf{U} &= \text{eigenvectors of } \mathbf{A}\mathbf{A}^T & n \times n \\ \mathbf{D} &= \sqrt{\text{diag}(\text{eig}(\mathbf{A}\mathbf{A}^T))} & n \times m \\ \mathbf{V} &= \text{eigenvectors of } \mathbf{A}^T\mathbf{A} & m \times m\end{aligned}$$

5.2.1 Symmetric Square decomposed into squares

Assume \mathbf{A} to be $n \times n$ and symmetric. Then

$$[\mathbf{A}] = [\mathbf{V}] [\mathbf{D}] [\mathbf{V}^T]$$

where \mathbf{D} is diagonal with the eigenvalues of \mathbf{A} and \mathbf{V} is orthogonal and the eigenvectors of \mathbf{A} .

5.2.2 Square decomposed into squares

Assume \mathbf{A} to be $n \times n$. Then

$$[\mathbf{A}] = [\mathbf{V}][\mathbf{D}][\mathbf{U}^T]$$

where \mathbf{D} is diagonal with the square root of the eigenvalues of $\mathbf{A}\mathbf{A}^T$, \mathbf{V} is the eigenvectors of $\mathbf{A}\mathbf{A}^T$ and \mathbf{U}^T is the eigenvectors of $\mathbf{A}^T\mathbf{A}$.

5.2.3 Square decomposed into rectangular

Assume $\mathbf{V}_*\mathbf{D}_*\mathbf{U}_*^T = \mathbf{0}$ then we can expand the SVD of \mathbf{A} into

$$[\mathbf{A}] = [\mathbf{V} \mid \mathbf{V}_*] \left[\begin{array}{c|c} \mathbf{D} & \mathbf{0} \\ \hline \mathbf{0} & \mathbf{D}_* \end{array} \right] \left[\begin{array}{c} \mathbf{U}^T \\ \hline \mathbf{U}_*^T \end{array} \right]$$

where the SVD of \mathbf{A} is $\mathbf{A} = \mathbf{V}\mathbf{D}\mathbf{U}^T$.

5.2.4 Rectangular decomposition I

Assume \mathbf{A} is $n \times m$

$$[\mathbf{A}] = [\mathbf{V}][\mathbf{D}][\mathbf{U}^T]$$

where \mathbf{D} is diagonal with the square root of the eigenvalues of $\mathbf{A}\mathbf{A}^T$, \mathbf{V} is the eigenvectors of $\mathbf{A}\mathbf{A}^T$ and \mathbf{U}^T is the eigenvectors of $\mathbf{A}^T\mathbf{A}$.

5.2.5 Rectangular decomposition II

Assume \mathbf{A} is $n \times m$

$$[\mathbf{A}] = [\mathbf{V}] \left[\begin{array}{c} \mathbf{D} \\ \hline \mathbf{0} \end{array} \right] \left[\begin{array}{c} \mathbf{U}^T \\ \hline \mathbf{0} \end{array} \right]$$

5.2.6 Rectangular decomposition III

Assume \mathbf{A} is $n \times m$

$$[\mathbf{A}] = [\mathbf{V}][\mathbf{D}] \left[\begin{array}{c} \mathbf{U}^T \\ \hline \mathbf{0} \end{array} \right]$$

where \mathbf{D} is diagonal with the square root of the eigenvalues of $\mathbf{A}\mathbf{A}^T$, \mathbf{V} is the eigenvectors of $\mathbf{A}\mathbf{A}^T$ and \mathbf{U}^T is the eigenvectors of $\mathbf{A}^T\mathbf{A}$.

5.3 Triangular Decomposition**5.3.1 Cholesky-decomposition**

Assume \mathbf{A} is positive definite, then

$$\mathbf{A} = \mathbf{B}^T\mathbf{B}$$

where \mathbf{B} is a unique upper triangular matrix.

6 General Statistics and Probability

6.1 Moments of any distribution

6.1.1 Mean and covariance of linear forms

Assume \mathbf{X} and \mathbf{x} to be a matrix and a vector of random variables. Then

$$E[\mathbf{AXB} + \mathbf{C}] = \mathbf{A}E[\mathbf{X}]\mathbf{B} + \mathbf{C}$$

$$\text{Var}[\mathbf{Ax}] = \mathbf{A}\text{Var}[\mathbf{x}]\mathbf{A}^T$$

$$\text{Cov}[\mathbf{Ax}, \mathbf{By}] = \mathbf{A}\text{Cov}[\mathbf{x}, \mathbf{y}]\mathbf{B}^T$$

See [14].

6.1.2 Mean and Variance of Square Forms

Assume \mathbf{A} is symmetric, $\mathbf{c} = E[\mathbf{x}]$ and $\mathbf{\Sigma} = \text{Var}[\mathbf{x}]$. Assume also that all coordinates x_i are independent, have the same central moments $\mu_1, \mu_2, \mu_3, \mu_4$ and denote $\mathbf{a} = \text{diag}(\mathbf{A})$. Then

$$E[\mathbf{x}^T \mathbf{Ax}] = \text{Tr}(\mathbf{A}\mathbf{\Sigma}) + \mathbf{c}^T \mathbf{Ac}$$

$$\text{Var}[\mathbf{x}^T \mathbf{Ax}] = 2\mu_2^2 \text{Tr}(\mathbf{A}^2) + 4\mu_2 \mathbf{c}^T \mathbf{A}^2 \mathbf{c} + 4\mu_3 \mathbf{c}^T \mathbf{Aa} + (\mu_4 - 3\mu_2^2) \mathbf{a}^T \mathbf{a}$$

See [14]

6.2 Expectations

Assume \mathbf{x} to be a stochastic vector with mean \mathbf{m} , covariance \mathbf{M} and central moments $\mathbf{v}_r = E[(\mathbf{x} - \mathbf{m})^r]$.

6.2.1 Linear Forms

$$E[\mathbf{Ax} + \mathbf{b}] = \mathbf{Am} + \mathbf{b}$$

$$E[\mathbf{Ax}] = \mathbf{Am}$$

$$E[\mathbf{x} + \mathbf{b}] = \mathbf{m} + \mathbf{b}$$

6.2.2 Quadratic Forms

$$E[(\mathbf{Ax} + \mathbf{a})(\mathbf{Bx} + \mathbf{b})^T] = \mathbf{AMB}^T + (\mathbf{Am} + \mathbf{a})(\mathbf{Bm} + \mathbf{b})^T$$

$$E[\mathbf{xx}^T] = \mathbf{M} + \mathbf{mm}^T$$

$$E[\mathbf{xa}^T \mathbf{x}] = (\mathbf{M} + \mathbf{mm}^T) \mathbf{a}$$

$$E[\mathbf{x}^T \mathbf{ax}^T] = \mathbf{a}^T (\mathbf{M} + \mathbf{mm}^T)$$

$$E[(\mathbf{Ax})(\mathbf{Ax})^T] = \mathbf{A}(\mathbf{M} + \mathbf{mm}^T) \mathbf{A}^T$$

$$E[(\mathbf{x} + \mathbf{a})(\mathbf{x} + \mathbf{a})^T] = \mathbf{M} + (\mathbf{m} + \mathbf{a})(\mathbf{m} + \mathbf{a})^T$$

$$\begin{aligned}
E[(\mathbf{Ax} + \mathbf{a})^T(\mathbf{Bx} + \mathbf{b})] &= \text{Tr}(\mathbf{AMB}^T) + (\mathbf{Am} + \mathbf{a})^T(\mathbf{Bm} + \mathbf{b}) \\
E[\mathbf{x}^T \mathbf{x}] &= \text{Tr}(\mathbf{M}) + \mathbf{m}^T \mathbf{m} \\
E[\mathbf{x}^T \mathbf{Ax}] &= \text{Tr}(\mathbf{AM}) + \mathbf{m}^T \mathbf{Am} \\
E[(\mathbf{Ax})^T(\mathbf{Ax})] &= \text{Tr}(\mathbf{AMA}^T) + (\mathbf{Am})^T(\mathbf{Am}) \\
E[(\mathbf{x} + \mathbf{a})^T(\mathbf{x} + \mathbf{a})] &= \text{Tr}(\mathbf{M}) + (\mathbf{m} + \mathbf{a})^T(\mathbf{m} + \mathbf{a})
\end{aligned}$$

See [7].

6.2.3 Cubic Forms

Assume \mathbf{x} to be independent, then

$$\begin{aligned}
E[(\mathbf{Ax} + \mathbf{a})(\mathbf{Bx} + \mathbf{b})^T(\mathbf{Cx} + \mathbf{c})] &= \mathbf{A} \text{diag}(\mathbf{B}^T \mathbf{C}) \mathbf{v}_3 \\
&\quad + \text{Tr}(\mathbf{BMC}^T)(\mathbf{Am} + \mathbf{a}) \\
&\quad + \mathbf{AMC}^T(\mathbf{Bm} + \mathbf{b}) \\
&\quad + (\mathbf{AMB}^T + (\mathbf{Am} + \mathbf{a})(\mathbf{Bm} + \mathbf{b})^T)(\mathbf{Cm} + \mathbf{c}) \\
E[\mathbf{xx}^T \mathbf{x}] &= \mathbf{v}_3 + 2\mathbf{Mm} + (\text{Tr}(\mathbf{M}) + \mathbf{m}^T \mathbf{m})\mathbf{m} \\
E[(\mathbf{Ax} + \mathbf{a})(\mathbf{Ax} + \mathbf{a})^T(\mathbf{Ax} + \mathbf{a})] &= \mathbf{A} \text{diag}(\mathbf{A}^T \mathbf{A}) \mathbf{v}_3 \\
&\quad + [2\mathbf{AMA}^T + (\mathbf{Ax} + \mathbf{a})(\mathbf{Ax} + \mathbf{a})^T](\mathbf{Am} + \mathbf{a}) \\
&\quad + \text{Tr}(\mathbf{AMA}^T)(\mathbf{Am} + \mathbf{a}) \\
E[(\mathbf{Ax} + \mathbf{a})\mathbf{b}^T(\mathbf{Cx} + \mathbf{c})(\mathbf{Dx} + \mathbf{d})^T] &= (\mathbf{Ax} + \mathbf{a})\mathbf{b}^T(\mathbf{CMD}^T + (\mathbf{Cm} + \mathbf{c})(\mathbf{Dm} + \mathbf{d})^T) \\
&\quad + (\mathbf{AMC}^T + (\mathbf{Am} + \mathbf{a})(\mathbf{Cm} + \mathbf{c})^T)\mathbf{b}(\mathbf{Dm} + \mathbf{d})^T \\
&\quad + \mathbf{b}^T(\mathbf{Cm} + \mathbf{c})(\mathbf{AMD}^T - (\mathbf{Am} + \mathbf{a})(\mathbf{Dm} + \mathbf{d})^T)
\end{aligned}$$

See [7].

7 Gaussians

7.1 One Dimensional

7.1.1 Density and Normalization

The density is

$$p(s) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{(s-\mu)^2}{2\sigma^2}\right)$$

Normalization integrals

$$\int e^{-\frac{(s-\mu)^2}{2\sigma^2}} ds = \sqrt{2\pi\sigma^2}$$

$$\int e^{-(ax^2+bx+c)} dx = \sqrt{\frac{\pi}{a}} \exp\left[\frac{b^2-4ac}{4a}\right]$$

$$\int e^{c_2x^2+c_1x+c_0} dx = \sqrt{\frac{\pi}{-c_2}} \exp\left[\frac{c_1^2-4c_2c_0}{-4c_2}\right]$$

7.1.2 Completing the Squares

$$c_2x^2 + c_1x + c_0 = -a(x-b)^2 + w$$

$$-a = c_2 \quad b = \frac{1}{2} \frac{c_1}{c_2} \quad w = \frac{1}{4} \frac{c_1^2}{c_2} + c_0$$

or

$$c_2x^2 + c_1x + c_0 = -\frac{1}{2\sigma^2}(x-\mu)^2 + d$$

$$\mu = \frac{-c_1}{2c_2} \quad \sigma^2 = \frac{-1}{2c_2} \quad d = c_0 - \frac{c_1^2}{4c_2}$$

7.1.3 Moments

If the density is expressed by

$$p(x) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left[-\frac{(s-\mu)^2}{2\sigma^2}\right] \quad \text{or} \quad p(x) = C \exp(c_2x^2 + c_1x)$$

then the first few basic moments are

$$\begin{aligned} \langle x \rangle &= \mu &= \frac{-c_1}{2c_2} \\ \langle x^2 \rangle &= \sigma^2 + \mu^2 &= \frac{-1}{2c_2} + \left(\frac{-c_1}{2c_2}\right)^2 \\ \langle x^3 \rangle &= 3\sigma^2\mu + \mu^3 &= \frac{c_1}{(2c_2)^2} \left[3 - \frac{c_1^2}{2c_2}\right] \\ \langle x^4 \rangle &= \mu^4 + 6\mu^2\sigma^2 + 3\sigma^4 &= \left(\frac{c_1}{2c_2}\right)^4 + 6\left(\frac{c_1}{2c_2}\right)^2 \left(\frac{-1}{2c_2}\right) + 3\left(\frac{1}{2c_2}\right)^2 \end{aligned}$$

and the central moments are

$$\begin{aligned}
\langle (x - \mu) \rangle &= 0 &= 0 \\
\langle (x - \mu)^2 \rangle &= \sigma^2 &= \left[\frac{-1}{2c_2} \right] \\
\langle (x - \mu)^3 \rangle &= 0 &= 0 \\
\langle (x - \mu)^4 \rangle &= 3\sigma^4 &= 3 \left[\frac{1}{2c_2} \right]^2
\end{aligned}$$

A kind of pseudo-moments (un-normalized integrals) can easily be derived as

$$\int \exp(c_2 x^2 + c_1 x) x^n dx = Z \langle x^n \rangle = \sqrt{\frac{\pi}{-c_2}} \exp \left[\frac{c_1^2}{-4c_2} \right] \langle x^n \rangle$$

From the un-centralized moments one can derive other entities like

$$\begin{aligned}
\langle x^2 \rangle - \langle x \rangle^2 &= \sigma^2 &= \frac{-1}{2c_2} \\
\langle x^3 \rangle - \langle x^2 \rangle \langle x \rangle &= 2\sigma^2 \mu &= \frac{2c_1}{(2c_2)^2} \\
\langle x^4 \rangle - \langle x^2 \rangle^2 &= 2\sigma^4 + 4\mu^2 \sigma^2 &= \frac{2}{(2c_2)^2} \left[1 - 4 \frac{c_1^2}{2c_2} \right]
\end{aligned}$$

7.2 Basics

7.2.1 Density and normalization

The density of $\mathbf{x} \sim \mathcal{N}(\mathbf{m}, \Sigma)$ is

$$p(\mathbf{x}) = \frac{1}{\sqrt{\det(2\pi\Sigma)}} \exp \left[-\frac{1}{2} (\mathbf{x} - \mathbf{m})^T \Sigma^{-1} (\mathbf{x} - \mathbf{m}) \right]$$

Note that if \mathbf{x} is d -dimensional, then $\det(2\pi\Sigma) = (2\pi)^d \det(\Sigma)$.

Integration and normalization

$$\begin{aligned}
\int \exp \left[-\frac{1}{2} (\mathbf{x} - \mathbf{m})^T \Sigma^{-1} (\mathbf{x} - \mathbf{m}) \right] d\mathbf{x} &= \sqrt{\det(2\pi\Sigma)} \\
\int \exp \left[-\frac{1}{2} \mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} \right] d\mathbf{x} &= \sqrt{\det(2\pi\mathbf{A}^{-1})} \exp \left[\frac{1}{2} \mathbf{b}^T \mathbf{A}^{-1} \mathbf{b} \right] \\
\int \exp \left[-\frac{1}{2} \text{Tr}(\mathbf{S}^T \mathbf{A} \mathbf{S}) + \text{Tr}(\mathbf{B}^T \mathbf{S}) \right] d\mathbf{S} &= \sqrt{\det(2\pi\mathbf{A}^{-1})} \exp \left[\frac{1}{2} \text{Tr}(\mathbf{B}^T \mathbf{A}^{-1} \mathbf{B}) \right]
\end{aligned}$$

The derivatives of the density are

$$\begin{aligned}
\frac{\partial p(\mathbf{x})}{\partial \mathbf{x}} &= -p(\mathbf{x}) \Sigma^{-1} (\mathbf{x} - \mathbf{m}) \\
\frac{\partial^2 p}{\partial \mathbf{x} \partial \mathbf{x}^T} &= p(\mathbf{x}) \left(\Sigma^{-1} (\mathbf{x} - \mathbf{m}) (\mathbf{x} - \mathbf{m})^T \Sigma^{-1} - \Sigma^{-1} \right)
\end{aligned}$$

7.2.2 Linear combination

Assume $\mathbf{x} \sim \mathcal{N}(\mathbf{m}_x, \Sigma_x)$ and $\mathbf{y} \sim \mathcal{N}(\mathbf{m}_y, \Sigma_y)$ then

$$\mathbf{A}\mathbf{x} + \mathbf{B}\mathbf{y} + \mathbf{c} \sim \mathcal{N}(\mathbf{A}\mathbf{m}_x + \mathbf{B}\mathbf{m}_y + \mathbf{c}, \mathbf{A}\Sigma_x\mathbf{A}^T + \mathbf{B}\Sigma_y\mathbf{B}^T)$$

7.2.3 Rearranging Means

$$\mathcal{N}_{\mathbf{Ax}}[\mathbf{m}, \boldsymbol{\Sigma}] = \frac{\sqrt{\det(2\pi(\mathbf{A}^T \boldsymbol{\Sigma}^{-1} \mathbf{A})^{-1})}}{\sqrt{\det(2\pi \boldsymbol{\Sigma})}} \mathcal{N}_{\mathbf{x}}[\mathbf{A}^{-1} \mathbf{m}, (\mathbf{A}^T \boldsymbol{\Sigma}^{-1} \mathbf{A})^{-1}]$$

7.2.4 Rearranging into squared form

If \mathbf{A} is symmetric, then

$$\begin{aligned} -\frac{1}{2} \mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} &= -\frac{1}{2} (\mathbf{x} - \mathbf{A}^{-1} \mathbf{b})^T \mathbf{A} (\mathbf{x} - \mathbf{A}^{-1} \mathbf{b}) + \frac{1}{2} \mathbf{b}^T \mathbf{A}^{-1} \mathbf{b} \\ -\frac{1}{2} \text{Tr}(\mathbf{X}^T \mathbf{A} \mathbf{X}) + \text{Tr}(\mathbf{B}^T \mathbf{X}) &= -\frac{1}{2} \text{Tr}[(\mathbf{X} - \mathbf{A}^{-1} \mathbf{B})^T \mathbf{A} (\mathbf{X} - \mathbf{A}^{-1} \mathbf{B})] + \frac{1}{2} \text{Tr}(\mathbf{B}^T \mathbf{A}^{-1} \mathbf{B}) \end{aligned}$$

7.2.5 Sum of two squared forms

In vector formulation (assuming $\boldsymbol{\Sigma}_1, \boldsymbol{\Sigma}_2$ are symmetric)

$$\begin{aligned} &-\frac{1}{2} (\mathbf{x} - \mathbf{m}_1)^T \boldsymbol{\Sigma}_1^{-1} (\mathbf{x} - \mathbf{m}_1) \\ &-\frac{1}{2} (\mathbf{x} - \mathbf{m}_2)^T \boldsymbol{\Sigma}_2^{-1} (\mathbf{x} - \mathbf{m}_2) \\ &= -\frac{1}{2} (\mathbf{x} - \mathbf{m}_c)^T \boldsymbol{\Sigma}_c^{-1} (\mathbf{x} - \mathbf{m}_c) + C \end{aligned}$$

$$\begin{aligned} \boldsymbol{\Sigma}_c^{-1} &= \boldsymbol{\Sigma}_1^{-1} + \boldsymbol{\Sigma}_2^{-1} \\ \mathbf{m}_c &= (\boldsymbol{\Sigma}_1^{-1} + \boldsymbol{\Sigma}_2^{-1})^{-1} (\boldsymbol{\Sigma}_1^{-1} \mathbf{m}_1 + \boldsymbol{\Sigma}_2^{-1} \mathbf{m}_2) \\ C &= \frac{1}{2} (\mathbf{m}_1^T \boldsymbol{\Sigma}_1^{-1} + \mathbf{m}_2^T \boldsymbol{\Sigma}_2^{-1}) (\boldsymbol{\Sigma}_1^{-1} + \boldsymbol{\Sigma}_2^{-1})^{-1} (\boldsymbol{\Sigma}_1^{-1} \mathbf{m}_1 + \boldsymbol{\Sigma}_2^{-1} \mathbf{m}_2) \\ &\quad - \frac{1}{2} (\mathbf{m}_1^T \boldsymbol{\Sigma}_1^{-1} \mathbf{m}_1 + \mathbf{m}_2^T \boldsymbol{\Sigma}_2^{-1} \mathbf{m}_2) \end{aligned}$$

In a trace formulation (assuming $\boldsymbol{\Sigma}_1, \boldsymbol{\Sigma}_2$ are symmetric)

$$\begin{aligned} &-\frac{1}{2} \text{Tr}((\mathbf{X} - \mathbf{M}_1)^T \boldsymbol{\Sigma}_1^{-1} (\mathbf{X} - \mathbf{M}_1)) \\ &-\frac{1}{2} \text{Tr}((\mathbf{X} - \mathbf{M}_2)^T \boldsymbol{\Sigma}_2^{-1} (\mathbf{X} - \mathbf{M}_2)) \\ &= -\frac{1}{2} \text{Tr}[(\mathbf{X} - \mathbf{M}_c)^T \boldsymbol{\Sigma}_c^{-1} (\mathbf{X} - \mathbf{M}_c)] + C \end{aligned}$$

$$\begin{aligned} \boldsymbol{\Sigma}_c^{-1} &= \boldsymbol{\Sigma}_1^{-1} + \boldsymbol{\Sigma}_2^{-1} \\ \mathbf{M}_c &= (\boldsymbol{\Sigma}_1^{-1} + \boldsymbol{\Sigma}_2^{-1})^{-1} (\boldsymbol{\Sigma}_1^{-1} \mathbf{M}_1 + \boldsymbol{\Sigma}_2^{-1} \mathbf{M}_2) \\ C &= \frac{1}{2} \text{Tr}[(\boldsymbol{\Sigma}_1^{-1} \mathbf{M}_1 + \boldsymbol{\Sigma}_2^{-1} \mathbf{M}_2)^T (\boldsymbol{\Sigma}_1^{-1} + \boldsymbol{\Sigma}_2^{-1})^{-1} (\boldsymbol{\Sigma}_1^{-1} \mathbf{M}_1 + \boldsymbol{\Sigma}_2^{-1} \mathbf{M}_2)] \\ &\quad - \frac{1}{2} \text{Tr}(\mathbf{M}_1^T \boldsymbol{\Sigma}_1^{-1} \mathbf{M}_1 + \mathbf{M}_2^T \boldsymbol{\Sigma}_2^{-1} \mathbf{M}_2) \end{aligned}$$

7.2.6 Product of gaussian densities

Let $\mathcal{N}_{\mathbf{x}}(\mathbf{m}, \mathbf{\Sigma})$ denote a density of \mathbf{x} , then

$$\mathcal{N}_{\mathbf{x}}(\mathbf{m}_1, \mathbf{\Sigma}_1) \cdot \mathcal{N}_{\mathbf{x}}(\mathbf{m}_2, \mathbf{\Sigma}_2) = c_c \mathcal{N}_{\mathbf{x}}(\mathbf{m}_c, \mathbf{\Sigma}_c)$$

$$\begin{aligned} c_c &= \mathcal{N}_{\mathbf{m}_1}(\mathbf{m}_2, (\mathbf{\Sigma}_1 + \mathbf{\Sigma}_2)) \\ &= \frac{1}{\sqrt{\det(2\pi(\mathbf{\Sigma}_1 + \mathbf{\Sigma}_2))}} \exp \left[-\frac{1}{2}(\mathbf{m}_1 - \mathbf{m}_2)^T (\mathbf{\Sigma}_1 + \mathbf{\Sigma}_2)^{-1} (\mathbf{m}_1 - \mathbf{m}_2) \right] \\ \mathbf{m}_c &= (\mathbf{\Sigma}_1^{-1} + \mathbf{\Sigma}_2^{-1})^{-1} (\mathbf{\Sigma}_1^{-1} \mathbf{m}_1 + \mathbf{\Sigma}_2^{-1} \mathbf{m}_2) \\ \mathbf{\Sigma}_c &= (\mathbf{\Sigma}_1^{-1} + \mathbf{\Sigma}_2^{-1})^{-1} \end{aligned}$$

but note that the product is not normalized as a density of \mathbf{x} .

7.3 Moments

7.3.1 Mean and covariance of linear forms

First and second moments. Assume $\mathbf{x} \sim \mathcal{N}(\mathbf{m}, \mathbf{\Sigma})$

$$E(\mathbf{x}) = \mathbf{m}$$

$$\text{Cov}(\mathbf{x}, \mathbf{x}) = \text{Var}(\mathbf{x}) = \mathbf{\Sigma} = E(\mathbf{x}\mathbf{x}^T) - E(\mathbf{x})E(\mathbf{x}^T) = E(\mathbf{x}\mathbf{x}^T) - \mathbf{m}\mathbf{m}^T$$

As for any other distribution it holds for gaussians that

$$E[\mathbf{A}\mathbf{x}] = \mathbf{A}E[\mathbf{x}]$$

$$\text{Var}[\mathbf{A}\mathbf{x}] = \mathbf{A}\text{Var}[\mathbf{x}]\mathbf{A}^T$$

$$\text{Cov}[\mathbf{A}\mathbf{x}, \mathbf{B}\mathbf{y}] = \mathbf{A}\text{Cov}[\mathbf{x}, \mathbf{y}]\mathbf{B}^T$$

7.3.2 Mean and variance of square forms

Mean and variance of square forms: Assume $\mathbf{x} \sim \mathcal{N}(\mathbf{m}, \mathbf{\Sigma})$

$$\begin{aligned} E(\mathbf{x}\mathbf{x}^T) &= \mathbf{\Sigma} + \mathbf{m}\mathbf{m}^T \\ E[\mathbf{x}^T \mathbf{A}\mathbf{x}] &= \text{Tr}(\mathbf{A}\mathbf{\Sigma}) + \mathbf{m}^T \mathbf{A}\mathbf{m} \\ \text{Var}(\mathbf{x}^T \mathbf{A}\mathbf{x}) &= 2\sigma^4 \text{Tr}(\mathbf{A}^2) + 4\sigma^2 \mathbf{m}^T \mathbf{A}^2 \mathbf{m} \\ E[(\mathbf{x} - \mathbf{m}')^T \mathbf{A}(\mathbf{x} - \mathbf{m}')] &= (\mathbf{m} - \mathbf{m}')^T \mathbf{A}(\mathbf{m} - \mathbf{m}') + \text{Tr}(\mathbf{A}\mathbf{\Sigma}) \end{aligned}$$

Assume $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I})$ and \mathbf{A} and \mathbf{B} to be symmetric, then

$$\text{Cov}(\mathbf{x}^T \mathbf{A}\mathbf{x}, \mathbf{x}^T \mathbf{B}\mathbf{x}) = 2\sigma^4 \text{Tr}(\mathbf{A}\mathbf{B})$$

7.3.3 Cubic forms

$$E[\mathbf{x}\mathbf{b}^T\mathbf{x}\mathbf{x}^T] = \mathbf{m}\mathbf{b}^T(\mathbf{M} + \mathbf{m}\mathbf{m}^T) + (\mathbf{M} + \mathbf{m}\mathbf{m}^T)\mathbf{b}\mathbf{m}^T + \mathbf{b}^T\mathbf{m}(\mathbf{M} - \mathbf{m}\mathbf{m}^T)$$

7.3.4 Mean of Quartic Forms

$$\begin{aligned} E[\mathbf{x}\mathbf{x}^T\mathbf{x}\mathbf{x}^T] &= 2(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)^2 + \mathbf{m}^T\mathbf{m}(\boldsymbol{\Sigma} - \mathbf{m}\mathbf{m}^T) \\ &\quad + \text{Tr}(\boldsymbol{\Sigma})(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T) \\ E[\mathbf{x}\mathbf{x}^T\mathbf{A}\mathbf{x}\mathbf{x}^T] &= (\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)(\mathbf{A} + \mathbf{A}^T)(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T) \\ &\quad + \mathbf{m}^T\mathbf{A}\mathbf{m}(\boldsymbol{\Sigma} - \mathbf{m}\mathbf{m}^T) + \text{Tr}[\mathbf{A}\boldsymbol{\Sigma}(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)] \\ E[\mathbf{x}^T\mathbf{x}\mathbf{x}^T\mathbf{x}] &= 2\text{Tr}(\boldsymbol{\Sigma}^2) + 4\mathbf{m}^T\boldsymbol{\Sigma}\mathbf{m} + (\text{Tr}(\boldsymbol{\Sigma}) + \mathbf{m}^T\mathbf{m})^2 \\ E[\mathbf{x}^T\mathbf{A}\mathbf{x}\mathbf{x}^T\mathbf{B}\mathbf{x}] &= \text{Tr}[\mathbf{A}\boldsymbol{\Sigma}(\mathbf{B} + \mathbf{B}^T)\boldsymbol{\Sigma}] + \mathbf{m}^T(\mathbf{A} + \mathbf{A}^T)\boldsymbol{\Sigma}(\mathbf{B} + \mathbf{B}^T)\mathbf{m} \\ &\quad + (\text{Tr}(\mathbf{A}\boldsymbol{\Sigma}) + \mathbf{m}^T\mathbf{A}\mathbf{m})(\text{Tr}(\mathbf{B}\boldsymbol{\Sigma}) + \mathbf{m}^T\mathbf{B}\mathbf{m}) \end{aligned}$$

$$\begin{aligned} &E[\mathbf{a}^T\mathbf{x}\mathbf{b}^T\mathbf{x}\mathbf{c}^T\mathbf{x}\mathbf{d}^T\mathbf{x}] \\ &= (\mathbf{a}^T(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)\mathbf{b})(\mathbf{c}^T(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)\mathbf{d}) \\ &\quad + (\mathbf{a}^T(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)\mathbf{c})(\mathbf{b}^T(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)\mathbf{d}) \\ &\quad + (\mathbf{a}^T(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)\mathbf{d})(\mathbf{b}^T(\boldsymbol{\Sigma} + \mathbf{m}\mathbf{m}^T)\mathbf{c}) - 2\mathbf{a}^T\mathbf{m}\mathbf{b}^T\mathbf{m}\mathbf{c}^T\mathbf{m}\mathbf{d}^T\mathbf{m} \end{aligned}$$

$$\begin{aligned} &E[(\mathbf{A}\mathbf{x} + \mathbf{a})(\mathbf{B}\mathbf{x} + \mathbf{b})^T(\mathbf{C}\mathbf{x} + \mathbf{c})(\mathbf{D}\mathbf{x} + \mathbf{d})^T] \\ &= [\mathbf{A}\boldsymbol{\Sigma}\mathbf{B}^T + (\mathbf{A}\mathbf{m} + \mathbf{a})(\mathbf{B}\mathbf{m} + \mathbf{b})^T][\mathbf{C}\boldsymbol{\Sigma}\mathbf{D}^T + (\mathbf{C}\mathbf{m} + \mathbf{c})(\mathbf{D}\mathbf{m} + \mathbf{d})^T] \\ &\quad + [\mathbf{A}\boldsymbol{\Sigma}\mathbf{C}^T + (\mathbf{A}\mathbf{m} + \mathbf{a})(\mathbf{C}\mathbf{m} + \mathbf{c})^T][\mathbf{B}\boldsymbol{\Sigma}\mathbf{D}^T + (\mathbf{B}\mathbf{m} + \mathbf{b})(\mathbf{D}\mathbf{m} + \mathbf{d})^T] \\ &\quad + (\mathbf{B}\mathbf{m} + \mathbf{b})^T(\mathbf{C}\mathbf{m} + \mathbf{c})[\mathbf{A}\boldsymbol{\Sigma}\mathbf{D}^T - (\mathbf{A}\mathbf{m} + \mathbf{a})(\mathbf{D}\mathbf{m} + \mathbf{d})^T] \\ &\quad + \text{Tr}(\mathbf{B}\boldsymbol{\Sigma}\mathbf{C}^T)[\mathbf{A}\boldsymbol{\Sigma}\mathbf{D}^T + (\mathbf{A}\mathbf{m} + \mathbf{a})(\mathbf{D}\mathbf{m} + \mathbf{d})^T] \end{aligned}$$

$$\begin{aligned} &E[(\mathbf{A}\mathbf{x} + \mathbf{a})^T(\mathbf{B}\mathbf{x} + \mathbf{b})(\mathbf{C}\mathbf{x} + \mathbf{c})^T(\mathbf{D}\mathbf{x} + \mathbf{d})] \\ &= \text{Tr}[\mathbf{A}\boldsymbol{\Sigma}(\mathbf{C}^T\mathbf{D} + \mathbf{D}^T\mathbf{C})\boldsymbol{\Sigma}\mathbf{B}^T] \\ &\quad + [(\mathbf{A}\mathbf{m} + \mathbf{a})^T\mathbf{B} + (\mathbf{B}\mathbf{m} + \mathbf{b})^T\mathbf{A}]\boldsymbol{\Sigma}[\mathbf{C}^T(\mathbf{D}\mathbf{m} + \mathbf{d}) + \mathbf{D}^T(\mathbf{C}\mathbf{m} + \mathbf{c})] \\ &\quad + [\text{Tr}(\mathbf{A}\boldsymbol{\Sigma}\mathbf{B}^T) + (\mathbf{A}\mathbf{m} + \mathbf{a})^T(\mathbf{B}\mathbf{m} + \mathbf{b})][\text{Tr}(\mathbf{C}\boldsymbol{\Sigma}\mathbf{D}^T) + (\mathbf{C}\mathbf{m} + \mathbf{c})^T(\mathbf{D}\mathbf{m} + \mathbf{d})] \end{aligned}$$

See [7].

7.3.5 Moments

$$\begin{aligned} E[\mathbf{x}] &= \sum_k \rho_k \mathbf{m}_k \\ \text{Cov}(\mathbf{x}) &= \sum_k \sum_{k'} \rho_k \rho_{k'} (\boldsymbol{\Sigma}_k + \mathbf{m}_k \mathbf{m}_k^T - \mathbf{m}_k \mathbf{m}_{k'}^T) \end{aligned}$$

7.4 Miscellaneous

7.4.1 Whitening

Assume $\mathbf{x} \sim \mathcal{N}(\mathbf{m}, \Sigma)$ then

$$\mathbf{z} = \Sigma^{-1/2}(\mathbf{x} - \mathbf{m}) \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$$

Conversely having $\mathbf{z} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$ one can generate data $\mathbf{x} \sim \mathcal{N}(\mathbf{m}, \Sigma)$ by setting

$$\mathbf{x} = \Sigma^{1/2}\mathbf{z} + \mathbf{m} \sim \mathcal{N}(\mathbf{m}, \Sigma)$$

Note that $\Sigma^{1/2}$ means the matrix which fulfils $\Sigma^{1/2}\Sigma^{1/2} = \Sigma$, and that it exists and is unique since Σ is positive definite.

7.4.2 The Chi-Square connection

Assume $\mathbf{x} \sim \mathcal{N}(\mathbf{m}, \Sigma)$ and \mathbf{x} to be n dimensional, then

$$z = (\mathbf{x} - \mathbf{m})^T \Sigma^{-1}(\mathbf{x} - \mathbf{m}) \sim \chi_n^2$$

7.4.3 Entropy

Entropy of a D -dimensional gaussian

$$H(\mathbf{x}) = \int \mathcal{N}(\mathbf{m}, \Sigma) \ln \mathcal{N}(\mathbf{m}, \Sigma) d\mathbf{x} = -\ln \sqrt{\det(2\pi\Sigma)} - \frac{D}{2}$$

7.5 One Dimensional Mixture of Gaussians

7.5.1 Density and Normalization

$$p(s) = \sum_k^K \frac{\rho_k}{\sqrt{2\pi\sigma_k^2}} \exp\left[-\frac{1}{2} \frac{(s - \mu_k)^2}{\sigma_k^2}\right]$$

7.5.2 Moments

An useful fact of MoG, is that

$$\langle x^n \rangle = \sum_k \rho_k \langle x^n \rangle_k$$

where $\langle \cdot \rangle_k$ denotes average with respect to the k .th component. We can calculate the first four moments from the densities

$$p(x) = \sum_k \rho_k \frac{1}{\sqrt{2\pi\sigma_k^2}} \exp\left[-\frac{1}{2} \frac{(x - \mu_k)^2}{\sigma_k^2}\right]$$

$$p(x) = \sum_k \rho_k C_k \exp[c_{k2}x^2 + c_{k1}x]$$

as

$$\begin{aligned}
\langle x \rangle &= \sum_k \rho_k \mu_k &= \sum_k \rho_k \left[\frac{-c_{k1}}{2c_{k2}} \right] \\
\langle x^2 \rangle &= \sum_k \rho_k (\sigma_k^2 + \mu_k^2) &= \sum_k \rho_k \left[\frac{-1}{2c_{k2}} + \left(\frac{-c_{k1}}{2c_{k2}} \right)^2 \right] \\
\langle x^3 \rangle &= \sum_k \rho_k (3\sigma_k^2 \mu_k + \mu_k^3) &= \sum_k \rho_k \left[\frac{c_{k1}}{(2c_{k2})^2} \left[3 - \frac{c_{k1}^2}{2c_{k2}} \right] \right] \\
\langle x^4 \rangle &= \sum_k \rho_k (\mu_k^4 + 6\mu_k^2 \sigma_k^2 + 3\sigma_k^4) &= \sum_k \rho_k \left[\left(\frac{1}{2c_{k2}} \right)^2 \left[\left(\frac{c_{k1}}{2c_{k2}} \right)^2 - 6 \frac{c_{k1}^2}{2c_{k2}} + 3 \right] \right]
\end{aligned}$$

If all the gaussians are centered, i.e. $\mu_k = 0$ for all k , then

$$\begin{aligned}
\langle x \rangle &= 0 &= 0 \\
\langle x^2 \rangle &= \sum_k \rho_k \sigma_k^2 &= \sum_k \rho_k \left[\frac{-1}{2c_{k2}} \right] \\
\langle x^3 \rangle &= 0 &= 0 \\
\langle x^4 \rangle &= \sum_k \rho_k 3\sigma_k^4 &= \sum_k \rho_k 3 \left[\frac{-1}{2c_{k2}} \right]^2
\end{aligned}$$

From the un-centralized moments one can derive other entities like

$$\begin{aligned}
\langle x^2 \rangle - \langle x \rangle^2 &= \sum_{k,k'} \rho_k \rho_{k'} [\mu_k^2 + \sigma_k^2 - \mu_k \mu_{k'}] \\
\langle x^3 \rangle - \langle x^2 \rangle \langle x \rangle &= \sum_{k,k'} \rho_k \rho_{k'} [3\sigma_k^2 \mu_k + \mu_k^3 - (\sigma_k^2 + \mu_k^2) \mu_{k'}] \\
\langle x^4 \rangle - \langle x^2 \rangle^2 &= \sum_{k,k'} \rho_k \rho_{k'} [\mu_k^4 + 6\mu_k^2 \sigma_k^2 + 3\sigma_k^4 - (\sigma_k^2 + \mu_k^2)(\sigma_{k'}^2 + \mu_{k'}^2)]
\end{aligned}$$

7.6 Mixture of Gaussians

7.6.1 Density

The variable \mathbf{x} is distributed as a mixture of gaussians if it has the density

$$p(\mathbf{x}) = \sum_{k=1}^K \rho_k \frac{1}{\sqrt{\det(2\pi \Sigma_k)}} \exp \left[-\frac{1}{2} (\mathbf{x} - \mathbf{m}_k)^T \Sigma_k^{-1} (\mathbf{x} - \mathbf{m}_k) \right]$$

where ρ_k sum to 1 and the Σ_k all are positive definite.

8 Miscellaneous

8.1 Functions and Series

8.1.1 Finite Series

$$(\mathbf{X}^n - \mathbf{I})(\mathbf{X} - \mathbf{I})^{-1} = \mathbf{I} + \mathbf{X} + \mathbf{X}^2 + \dots + \mathbf{X}^{n-1}$$

8.1.2 Taylor Expansion of Scalar Function

Consider some scalar function $f(\mathbf{x})$ which takes the vector \mathbf{x} as an argument. This we can Taylor expand around \mathbf{x}_0

$$f(\mathbf{x}) \cong f(\mathbf{x}_0) + \mathbf{g}(\mathbf{x}_0)^T(\mathbf{x} - \mathbf{x}_0) + \frac{1}{2}(\mathbf{x} - \mathbf{x}_0)^T \mathbf{H}(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)$$

where

$$\mathbf{g}(\mathbf{x}_0) = \left. \frac{\partial f(\mathbf{x})}{\partial \mathbf{x}} \right|_{\mathbf{x}_0} \quad \mathbf{H}(\mathbf{x}_0) = \left. \frac{\partial^2 f(\mathbf{x})}{\partial \mathbf{x} \partial \mathbf{x}^T} \right|_{\mathbf{x}_0}$$

8.1.3 Taylor Expansion of Vector Functions

8.1.4 Matrix Functions by Infinite Series

As for analytical functions in one dimension, one can define a matrix function for square matrices \mathbf{X} by an infinite series

$$\mathbf{f}(\mathbf{X}) = \sum_{n=0}^{\infty} c_n \mathbf{X}^n$$

assuming the limit exists and is finite. If the coefficients c_n fulfils $\sum_n c_n x^n < \infty$, then one can prove that the above series exists and is finite, see [1]. Thus for any analytical function $f(x)$ there exists a corresponding matrix function $\mathbf{f}(\mathbf{x})$ constructed by the Taylor expansion. Using this one can prove the following results:

1) A matrix \mathbf{A} is a zero of its own characteristic polynomial [1]:

$$p(\lambda) = \det(\mathbf{I}\lambda - \mathbf{A}) = \sum_n c_n \lambda^n \quad \Rightarrow \quad p(\mathbf{A}) = \mathbf{0}$$

2) If \mathbf{A} is square it holds that [1]

$$\mathbf{A} = \mathbf{U}\mathbf{B}\mathbf{U}^{-1} \quad \Rightarrow \quad \mathbf{f}(\mathbf{A}) = \mathbf{U}\mathbf{f}(\mathbf{B})\mathbf{U}^{-1}$$

3) A useful fact when using power series is that

$$\mathbf{A}^n \rightarrow \mathbf{0} \text{ for } n \rightarrow \infty \quad \text{if} \quad |\mathbf{A}| < 1$$

8.1.5 Exponential Matrix Function

In analogy to the ordinary scalar exponential function, one can define exponential and logarithmic matrix functions:

$$e^{\mathbf{A}} = \sum_{n=0}^{\infty} \frac{1}{n!} \mathbf{A}^n = \mathbf{I} + \mathbf{A} + \frac{1}{2} \mathbf{A}^2 + \dots$$

$$e^{-\mathbf{A}} = \sum_{n=0}^{\infty} \frac{1}{n!} (-1)^n \mathbf{A}^n = \mathbf{I} - \mathbf{A} + \frac{1}{2} \mathbf{A}^2 - \dots$$

$$e^{t\mathbf{A}} = \sum_{n=0}^{\infty} \frac{1}{n!} (t\mathbf{A})^n = \mathbf{I} + t\mathbf{A} + \frac{1}{2} t^2 \mathbf{A}^2 + \dots$$

$$\ln(\mathbf{I} + \mathbf{A}) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}}{n} \mathbf{A}^n = \mathbf{A} - \frac{1}{2} \mathbf{A}^2 + \frac{1}{3} \mathbf{A}^3 - \dots$$

Some of the properties of the exponential function are [1]

$$e^{\mathbf{A}} e^{\mathbf{B}} = e^{\mathbf{A}+\mathbf{B}} \quad \text{if} \quad \mathbf{AB} = \mathbf{BA}$$

$$(e^{\mathbf{A}})^{-1} = e^{-\mathbf{A}}$$

$$\frac{d}{dt} e^{t\mathbf{A}} = \mathbf{A} e^{t\mathbf{A}}, \quad t \in \mathbb{R}$$

8.1.6 Trigonometric Functions

$$\sin(\mathbf{A}) = \sum_{n=0}^{\infty} \frac{(-1)^n \mathbf{A}^{2n+1}}{(2n+1)!} = \mathbf{A} - \frac{1}{3!} \mathbf{A}^3 + \frac{1}{5!} \mathbf{A}^5 - \dots$$

$$\cos(\mathbf{A}) = \sum_{n=0}^{\infty} \frac{(-1)^n \mathbf{A}^{2n}}{(2n)!} = \mathbf{I} - \frac{1}{2!} \mathbf{A}^2 + \frac{1}{4!} \mathbf{A}^4 - \dots$$

8.2 Indices, Entries and Vectors

Let \mathbf{e}_i denote the column vector which is 1 on entry i and zero elsewhere, i.e. $(\mathbf{e}_i)_j = \delta_{ij}$, and let \mathbf{J}^{ij} denote the matrix which is 1 on entry (i, j) and zero elsewhere.

8.2.1 Rows and Columns

$$i.\text{th row of } \mathbf{A} = \mathbf{e}_i^T \mathbf{A}$$

$$j.\text{th column of } \mathbf{A} = \mathbf{A} \mathbf{e}_j$$

8.2.2 Permutations

Let \mathbf{P} be some permutation matrix, e.g.

$$\mathbf{P} = \begin{bmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix} = [\mathbf{e}_2 \quad \mathbf{e}_1 \quad \mathbf{e}_3] = \begin{bmatrix} \mathbf{e}_2^T \\ \mathbf{e}_1^T \\ \mathbf{e}_3^T \end{bmatrix}$$

then

$$\mathbf{AP} = [\mathbf{Ae}_2 \quad \mathbf{Ae}_1 \quad \mathbf{Ae}_3] \quad \mathbf{PA} = \begin{bmatrix} \mathbf{e}_2^T \mathbf{A} \\ \mathbf{e}_1^T \mathbf{A} \\ \mathbf{e}_3^T \mathbf{A} \end{bmatrix}$$

That is, the first is a matrix which has columns of \mathbf{A} but in permuted sequence and the second is a matrix which has the rows of \mathbf{A} but in the permuted sequence.

8.2.3 Swap and Zeros

Assume \mathbf{A} to be $n \times m$ and \mathbf{J}^{ij} to be $m \times p$

$$\mathbf{AJ}^{ij} = [\mathbf{0} \quad \mathbf{0} \quad \dots \quad \mathbf{A}_i \quad \dots \quad \mathbf{0}]$$

i.e. an $n \times p$ matrix of zeros with the i .th column of \mathbf{A} in the place of the j .th column. Assume \mathbf{A} to be $n \times m$ and \mathbf{J}^{ij} to be $p \times n$

$$\mathbf{J}^{ij} \mathbf{A} = \begin{bmatrix} \mathbf{0} \\ \mathbf{0} \\ \vdots \\ \mathbf{A}_j \\ \vdots \\ \mathbf{0} \end{bmatrix}$$

i.e. a $p \times m$ matrix of zeros with the j .th row of \mathbf{A} in the place of the i .th row.

8.2.4 Rewriting product of elements

$$\begin{aligned} A_{ki} B_{jl} &= (\mathbf{Ae}_i \mathbf{e}_j^T \mathbf{B})_{kl} = (\mathbf{AJ}^{ij} \mathbf{B})_{kl} \\ A_{ik} B_{lj} &= (\mathbf{A}^T \mathbf{e}_i \mathbf{e}_j^T \mathbf{B}^T)_{kl} = (\mathbf{A}^T \mathbf{J}^{ij} \mathbf{B}^T)_{kl} \\ A_{ik} B_{jl} &= (\mathbf{A}^T \mathbf{e}_i \mathbf{e}_j^T \mathbf{B})_{kl} = (\mathbf{A}^T \mathbf{J}^{ij} \mathbf{B})_{kl} \\ A_{ki} B_{lj} &= (\mathbf{Ae}_i \mathbf{e}_j^T \mathbf{B}^T)_{kl} = (\mathbf{AJ}^{ij} \mathbf{B}^T)_{kl} \end{aligned}$$

8.2.5 Properties of the Singleentry MatrixIf $i = j$

$$\begin{aligned}\mathbf{J}^{ij} \mathbf{J}^{ij} &= \mathbf{J}^{ij} & (\mathbf{J}^{ij})^T (\mathbf{J}^{ij})^T &= \mathbf{J}^{ij} \\ \mathbf{J}^{ij} (\mathbf{J}^{ij})^T &= \mathbf{J}^{ij} & (\mathbf{J}^{ij})^T \mathbf{J}^{ij} &= \mathbf{J}^{ij}\end{aligned}$$

If $i \neq j$

$$\begin{aligned}\mathbf{J}^{ij} \mathbf{J}^{ij} &= \mathbf{0} & (\mathbf{J}^{ij})^T (\mathbf{J}^{ij})^T &= \mathbf{0} \\ \mathbf{J}^{ij} (\mathbf{J}^{ij})^T &= \mathbf{J}^{ii} & (\mathbf{J}^{ij})^T \mathbf{J}^{ij} &= \mathbf{J}^{jj}\end{aligned}$$

8.2.6 The Singleentry Matrix in Scalar ExpressionsAssume \mathbf{A} is $n \times m$ and \mathbf{J} is $m \times n$, then

$$\text{Tr}(\mathbf{A}\mathbf{J}^{ij}) = \text{Tr}(\mathbf{J}^{ij}\mathbf{A}) = (\mathbf{A}^T)_{ij}$$

Assume \mathbf{A} is $n \times n$, \mathbf{J} is $n \times m$ and \mathbf{B} is $m \times n$, then

$$\begin{aligned}\text{Tr}(\mathbf{A}\mathbf{J}^{ij}\mathbf{B}) &= (\mathbf{A}^T \mathbf{B}^T)_{ij} \\ \text{Tr}(\mathbf{A}\mathbf{J}^{ji}\mathbf{B}) &= (\mathbf{B}\mathbf{A})_{ij} \\ \text{Tr}(\mathbf{A}\mathbf{J}^{ij}\mathbf{J}^{ij}\mathbf{B}) &= \text{diag}(\mathbf{A}^T \mathbf{B}^T)_{ij}\end{aligned}$$

Assume \mathbf{A} is $n \times n$, \mathbf{J}^{ij} is $n \times m$ \mathbf{B} is $m \times n$, then

$$\begin{aligned}\mathbf{x}^T \mathbf{A}\mathbf{J}^{ij}\mathbf{B}\mathbf{x} &= (\mathbf{A}^T \mathbf{x}\mathbf{x}^T \mathbf{B}^T)_{ij} \\ \mathbf{x}^T \mathbf{A}\mathbf{J}^{ij}\mathbf{J}^{ij}\mathbf{B}\mathbf{x} &= \text{diag}(\mathbf{A}^T \mathbf{x}\mathbf{x}^T \mathbf{B}^T)_{ij}\end{aligned}$$

8.2.7 Structure Matrices

The structure matrix is defined by

$$\frac{\partial \mathbf{A}}{\partial A_{ij}} = \mathbf{S}^{ij}$$

If \mathbf{A} has no special structure then

$$\mathbf{S}^{ij} = \mathbf{J}^{ij}$$

If \mathbf{A} is symmetric then

$$\mathbf{S}^{ij} = \mathbf{J}^{ij} + \mathbf{J}^{ji} - \mathbf{J}^{ij}\mathbf{J}^{ij}$$

8.3 Solutions to Systems of Equations

8.3.1 Existence in Linear Systems

Assume \mathbf{A} is $n \times m$ and consider the linear system

$$\mathbf{Ax} = \mathbf{b}$$

Construct the augmented matrix $\mathbf{B} = [\mathbf{A} \ \mathbf{b}]$ then

<i>Condition</i>	<i>Solution</i>
$\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{B}) = m$	Unique solution \mathbf{x}
$\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{B}) < m$	Many solutions \mathbf{x}
$\text{rank}(\mathbf{A}) < \text{rank}(\mathbf{B})$	No solutions \mathbf{x}

8.3.2 Standard Square

Assume \mathbf{A} is square and invertible, then

$$\mathbf{Ax} = \mathbf{b} \quad \Rightarrow \quad \mathbf{x} = \mathbf{A}^{-1}\mathbf{b}$$

8.3.3 Degenerated Square

8.3.4 Over-determined Rectangular

Assume \mathbf{A} to be $n \times m$, $n > m$ (tall) and $\text{rank}(\mathbf{A}) = m$, then

$$\mathbf{Ax} = \mathbf{b} \quad \Rightarrow \quad \mathbf{x} = (\mathbf{A}^T \mathbf{A})^{-1} \mathbf{A}^T \mathbf{b} = \mathbf{A}^+ \mathbf{b}$$

that is *if* there exists a solution \mathbf{x} at all! If there is no solution the following can be useful:

$$\mathbf{Ax} = \mathbf{b} \quad \Rightarrow \quad \mathbf{x}_{min} = \mathbf{A}^+ \mathbf{b}$$

Now \mathbf{x}_{min} is the vector \mathbf{x} which minimizes $\|\mathbf{Ax} - \mathbf{b}\|^2$, i.e. the vector which is "least wrong". The matrix \mathbf{A}^+ is the pseudo-inverse of \mathbf{A} . See [3].

8.3.5 Under-determined Rectangular

Assume \mathbf{A} is $n \times m$ and $n < m$ ("broad").

$$\mathbf{Ax} = \mathbf{b} \quad \Rightarrow \quad \mathbf{x}_{min} = \mathbf{A}^T (\mathbf{A} \mathbf{A}^T)^{-1} \mathbf{b}$$

The equation have many solutions \mathbf{x} . But \mathbf{x}_{min} is the solution which minimizes $\|\mathbf{Ax} - \mathbf{b}\|^2$ and also the solution with the smallest norm $\|\mathbf{x}\|^2$. The same holds for a matrix version: Assume \mathbf{A} is $n \times m$, \mathbf{X} is $m \times n$ and \mathbf{B} is $n \times n$, then

$$\mathbf{AX} = \mathbf{B} \quad \Rightarrow \quad \mathbf{X}_{min} = \mathbf{A}^+ \mathbf{B}$$

The equation have many solutions \mathbf{X} . But \mathbf{X}_{min} is the solution which minimizes $\|\mathbf{AX} - \mathbf{B}\|^2$ and also the solution with the smallest norm $\|\mathbf{X}\|^2$. See [3].

Similar but different: Assume \mathbf{A} is square $n \times n$ and the matrices $\mathbf{B}_0, \mathbf{B}_1$ are $n \times N$, where $N > n$, then if \mathbf{B}_0 has maximal rank

$$\mathbf{A}\mathbf{B}_0 = \mathbf{B}_1 \quad \Rightarrow \quad \mathbf{A}_{min} = \mathbf{B}_1\mathbf{B}_0^T(\mathbf{B}_0\mathbf{B}_0^T)^{-1}$$

where \mathbf{A}_{min} denotes the matrix which is optimal in a least square sense. An interpretation is that \mathbf{A} is the linear approximation which maps the columns vectors of \mathbf{B}_0 into the columns vectors of \mathbf{B}_1 .

8.3.6 Linear form and zeros

$$\mathbf{A}\mathbf{x} = \mathbf{0}, \quad \forall \mathbf{x} \quad \Rightarrow \quad \mathbf{A} = \mathbf{0}$$

8.3.7 Square form and zeros

If \mathbf{A} is symmetric, then

$$\mathbf{x}^T \mathbf{A} \mathbf{x} = 0, \quad \forall \mathbf{x} \quad \Rightarrow \quad \mathbf{A} = \mathbf{0}$$

8.4 Block matrices

Let \mathbf{A}_{ij} denote the ij .th block of \mathbf{A} .

8.4.1 Multiplication

Assuming the dimensions of the blocks matches we have

$$\left[\begin{array}{c|c} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{array} \right] \left[\begin{array}{c|c} \mathbf{B}_{11} & \mathbf{B}_{12} \\ \mathbf{B}_{21} & \mathbf{B}_{22} \end{array} \right] = \left[\begin{array}{c|c} \mathbf{A}_{11}\mathbf{B}_{11} + \mathbf{A}_{12}\mathbf{B}_{21} & \mathbf{A}_{11}\mathbf{B}_{12} + \mathbf{A}_{12}\mathbf{B}_{22} \\ \mathbf{A}_{21}\mathbf{B}_{11} + \mathbf{A}_{22}\mathbf{B}_{21} & \mathbf{A}_{21}\mathbf{B}_{12} + \mathbf{A}_{22}\mathbf{B}_{22} \end{array} \right]$$

8.4.2 The Determinant

The determinant can be expressed as by the use of

$$\begin{aligned} \mathbf{C}_1 &= \mathbf{A}_{11} - \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21} \\ \mathbf{C}_2 &= \mathbf{A}_{22} - \mathbf{A}_{21}\mathbf{A}_{11}^{-1}\mathbf{A}_{12} \end{aligned}$$

as

$$\det \left(\left[\begin{array}{c|c} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{array} \right] \right) = \det(\mathbf{A}_{22}) \cdot \det(\mathbf{C}_1) = \det(\mathbf{A}_{11}) \cdot \det(\mathbf{C}_2)$$

8.4.3 The Inverse

The inverse can be expressed as by the use of

$$\begin{aligned} \mathbf{C}_1 &= \mathbf{A}_{11} - \mathbf{A}_{12}\mathbf{A}_{22}^{-1}\mathbf{A}_{21} \\ \mathbf{C}_2 &= \mathbf{A}_{22} - \mathbf{A}_{21}\mathbf{A}_{11}^{-1}\mathbf{A}_{12} \end{aligned}$$

as

$$\begin{aligned} \left[\begin{array}{c|c} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{A}_{21} & \mathbf{A}_{22} \end{array} \right]^{-1} &= \left[\begin{array}{c|c} \mathbf{C}_1^{-1} & -\mathbf{A}_{11}^{-1}\mathbf{A}_{12}\mathbf{C}_2^{-1} \\ -\mathbf{C}_2^{-1}\mathbf{A}_{21}\mathbf{A}_{11}^{-1} & \mathbf{C}_2^{-1} \end{array} \right] \\ &= \left[\begin{array}{c|c} \mathbf{A}_{11}^{-1} + \mathbf{A}_{11}^{-1}\mathbf{A}_{12}\mathbf{C}_2^{-1}\mathbf{A}_{21}\mathbf{A}_{11}^{-1} & -\mathbf{C}_1^{-1}\mathbf{A}_{12}\mathbf{A}_{22}^{-1} \\ -\mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{C}_1^{-1} & \mathbf{A}_{22}^{-1} + \mathbf{A}_{22}^{-1}\mathbf{A}_{21}\mathbf{C}_1^{-1}\mathbf{A}_{12}\mathbf{A}_{22}^{-1} \end{array} \right] \end{aligned}$$

8.4.4 Block diagonal

For block diagonal matrices we have

$$\begin{aligned} \left[\begin{array}{c|c} \mathbf{A}_{11} & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_{22} \end{array} \right]^{-1} &= \left[\begin{array}{c|c} (\mathbf{A}_{11})^{-1} & \mathbf{0} \\ \mathbf{0} & (\mathbf{A}_{22})^{-1} \end{array} \right] \\ \det \left(\left[\begin{array}{c|c} \mathbf{A}_{11} & \mathbf{0} \\ \mathbf{0} & \mathbf{A}_{22} \end{array} \right] \right) &= \det(\mathbf{A}_{11}) \cdot \det(\mathbf{A}_{22}) \end{aligned}$$

8.5 Matrix Norms

8.5.1 Definitions

A matrix norm is a mapping which fulfils

$$\begin{aligned} \|\mathbf{A}\| &\geq 0 & \|\mathbf{A}\| = 0 &\Leftrightarrow \mathbf{A} = \mathbf{0} \\ \|c\mathbf{A}\| &= |c|\|\mathbf{A}\|, & c &\in \mathbb{R} \\ \|\mathbf{A} + \mathbf{B}\| &\leq \|\mathbf{A}\| + \|\mathbf{B}\| \end{aligned}$$

8.5.2 Examples

$$\begin{aligned} \|\mathbf{A}\|_1 &= \max_j \sum_i |A_{ij}| \\ \|\mathbf{A}\|_2 &= \sqrt{\max \text{eig}(\mathbf{A}^T \mathbf{A})} \\ \|\mathbf{A}\|_p &= \left(\max_{\|\mathbf{x}\|_p=1} \|\mathbf{A}\mathbf{x}\|_p \right)^{1/p} \\ \|\mathbf{A}\|_\infty &= \max_i \sum_j |A_{ij}| \\ \|\mathbf{A}\|_F &= \sqrt{\sum_{ij} |A_{ij}|^2} \quad (\text{Frobenius}) \\ \|\mathbf{A}\|_{max} &= \max_{ij} |A_{ij}| \\ \|\mathbf{A}\|_{KF} &= \|\text{sing}(\mathbf{A})\|_1 \quad (\text{Ky Fan}) \end{aligned}$$

where $\text{sing}(\mathbf{A})$ is the vector of singular values of the matrix \mathbf{A} .

8.5.3 Inequalities

E. H. Rasmussen has in yet unpublished material derived and collected the following inequalities. They are collected in a table as below, assuming \mathbf{A} is an $m \times n$, and $d = \min\{m, n\}$

	$\ \mathbf{A}\ _{max}$	$\ \mathbf{A}\ _1$	$\ \mathbf{A}\ _\infty$	$\ \mathbf{A}\ _2$	$\ \mathbf{A}\ _F$	$\ \mathbf{A}\ _{KF}$
$\ \mathbf{A}\ _{max}$		1	1	1	1	1
$\ \mathbf{A}\ _1$	m		m	\sqrt{m}	\sqrt{m}	\sqrt{m}
$\ \mathbf{A}\ _\infty$	n	n		\sqrt{n}	\sqrt{n}	\sqrt{n}
$\ \mathbf{A}\ _2$	\sqrt{mn}	\sqrt{n}	\sqrt{m}		1	1
$\ \mathbf{A}\ _F$	\sqrt{mn}	\sqrt{n}	\sqrt{m}	\sqrt{d}		1
$\ \mathbf{A}\ _{KF}$	\sqrt{mnd}	\sqrt{nd}	\sqrt{md}	d	\sqrt{d}	

which are to be read as, e.g.

$$\|\mathbf{A}\|_2 \leq \sqrt{m} \cdot \|\mathbf{A}\|_\infty$$

8.6 Positive Definite and Semi-definite Matrices

8.6.1 Definitions

A matrix \mathbf{A} is positive definite if and only if

$$\mathbf{x}^T \mathbf{A} \mathbf{x} > \mathbf{0}, \quad \forall \mathbf{x}$$

A matrix \mathbf{A} is positive semi-definite if and only if

$$\mathbf{x}^T \mathbf{A} \mathbf{x} \geq \mathbf{0}, \quad \forall \mathbf{x}$$

Note that if \mathbf{A} is positive definite, then \mathbf{A} is also positive semi-definite.

8.6.2 Eigenvalues

The following holds with respect to the eigenvalues:

$$\begin{aligned} \mathbf{A} \text{ pos. def.} &\Rightarrow \text{eig}(\mathbf{A}) > 0 \\ \mathbf{A} \text{ pos. semi-def.} &\Rightarrow \text{eig}(\mathbf{A}) \geq 0 \end{aligned}$$

8.6.3 Trace

The following holds with respect to the trace:

$$\begin{aligned} \mathbf{A} \text{ pos. def.} &\Rightarrow \text{Tr}(\mathbf{A}) > 0 \\ \mathbf{A} \text{ pos. semi-def.} &\Rightarrow \text{Tr}(\mathbf{A}) \geq 0 \end{aligned}$$

8.6.4 Inverse

If \mathbf{A} is positive definite, then \mathbf{A} is invertible and \mathbf{A}^{-1} is also positive definite.

8.6.5 Diagonal

If \mathbf{A} is positive definite, then $A_{ii} > 0, \forall i$

8.6.6 Decomposition I

The matrix \mathbf{A} is positive semi-definite of rank $r \Leftrightarrow$ there exists a matrix \mathbf{B} of rank r such that $\mathbf{A} = \mathbf{B}\mathbf{B}^T$

The matrix \mathbf{A} is positive definite \Leftrightarrow there exists an invertible matrix \mathbf{B} such that $\mathbf{A} = \mathbf{B}\mathbf{B}^T$

8.6.7 Decomposition II

Assume \mathbf{A} is an $n \times n$ positive semi-definite, then there exists an $n \times r$ matrix \mathbf{B} of rank r such that $\mathbf{B}^T \mathbf{A} \mathbf{B} = \mathbf{I}$.

8.6.8 Equation with zeros

Assume \mathbf{A} is positive semi-definite, then $\mathbf{X}^T \mathbf{A} \mathbf{X} = \mathbf{0} \Rightarrow \mathbf{A} \mathbf{X} = \mathbf{0}$

8.6.9 Rank of product

Assume \mathbf{A} is positive definite, then $\text{rank}(\mathbf{B}\mathbf{A}\mathbf{B}^T) = \text{rank}(\mathbf{B})$

8.6.10 Positive definite property

If \mathbf{A} is $n \times n$ positive definite and \mathbf{B} is $r \times n$ of rank r , then $\mathbf{B}\mathbf{A}\mathbf{B}^T$ is positive definite.

8.6.11 Outer Product

If \mathbf{X} is $n \times r$ of rank r , then $\mathbf{X}\mathbf{X}^T$ is positive definite.

8.6.12 Small perturbations

If \mathbf{A} is positive definite and \mathbf{B} is symmetric, then $\mathbf{A} - t\mathbf{B}$ is positive definite for sufficiently small t .

8.7 Integral Involving Dirac Delta Functions

Assuming \mathbf{A} to be square, then

$$\int p(\mathbf{s})\delta(\mathbf{x} - \mathbf{A}\mathbf{s})d\mathbf{s} = \frac{1}{\det(\mathbf{A})}p(\mathbf{A}^{-1}\mathbf{x})$$

Assuming \mathbf{A} to be "underdetermined", i.e. "tall", then

$$\int p(\mathbf{s})\delta(\mathbf{x} - \mathbf{A}\mathbf{s})d\mathbf{s} = \begin{cases} \frac{1}{\sqrt{\det(\mathbf{A}^T \mathbf{A})}}p(\mathbf{A}^+ \mathbf{x}) & \text{if } \mathbf{x} = \mathbf{A}\mathbf{A}^+ \mathbf{x} \\ 0 & \text{elsewhere} \end{cases}$$

See [8].

8.8 Miscellaneous

For any \mathbf{A} it holds that

$$\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}^T) = \text{rank}(\mathbf{A}\mathbf{A}^T) = \text{rank}(\mathbf{A}^T\mathbf{A})$$

Assume \mathbf{A} is positive definite. Then

$$\text{rank}(\mathbf{B}^T\mathbf{A}\mathbf{B}) = \text{rank}(\mathbf{B})$$

$$\mathbf{A} \text{ is positive definite} \iff \exists \mathbf{B} \text{ invertible, such that } \mathbf{A} = \mathbf{B}\mathbf{B}^T$$

A Proofs and Details

A.0.1 Proof of Equation 14

Essentially we need to calculate

$$\begin{aligned}
 \frac{\partial(\mathbf{X}^n)_{kl}}{\partial X_{ij}} &= \frac{\partial}{\partial X_{ij}} \sum_{u_1, \dots, u_{n-1}} X_{k,u_1} X_{u_1,u_2} \dots X_{u_{n-1},l} \\
 &= \delta_{k,i} \delta_{u_1,j} X_{u_1,u_2} \dots X_{u_{n-1},l} \\
 &\quad + X_{k,u_1} \delta_{u_1,i} \delta_{u_2,j} \dots X_{u_{n-1},l} \\
 &\quad \vdots \\
 &\quad + X_{k,u_1} X_{u_1,u_2} \dots \delta_{u_{n-1},i} \delta_{l,j} \\
 &= \sum_{r=0}^{n-1} (\mathbf{X}^r)_{ki} (\mathbf{X}^{n-1-r})_{jl} \\
 &= \sum_{r=0}^{n-1} (\mathbf{X}^r \mathbf{J}^{ij} \mathbf{X}^{n-1-r})_{kl}
 \end{aligned}$$

Using the properties of the single entry matrix found in Sec. 8.2.5, the result follows easily.

A.0.2 Details on Eq. 47

$$\begin{aligned}
 \partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) \text{Tr}[(\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \partial(\mathbf{X}^H \mathbf{A} \mathbf{X})] \\
 &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) \text{Tr}[(\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} (\partial(\mathbf{X}^H) \mathbf{A} \mathbf{X} + \mathbf{X}^H \partial(\mathbf{A} \mathbf{X}))] \\
 &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) (\text{Tr}[(\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \partial(\mathbf{X}^H) \mathbf{A} \mathbf{X}] + \text{Tr}[(\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \mathbf{X}^H \partial(\mathbf{A} \mathbf{X})]) \\
 &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) (\text{Tr}[\mathbf{A} \mathbf{X} (\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \partial(\mathbf{X}^H)] + \text{Tr}[(\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \mathbf{X}^H \mathbf{A} \partial(\mathbf{X})])
 \end{aligned}$$

First, the derivative is found with respect to the real part of \mathbf{X}

$$\begin{aligned}
 \frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \Re \mathbf{X}} &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) \left(\frac{\text{Tr}[\mathbf{A} \mathbf{X} (\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \partial(\mathbf{X}^H)]}{\partial \Re \mathbf{X}} + \frac{\text{Tr}[(\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \mathbf{X}^H \mathbf{A} \partial(\mathbf{X})]}{\partial \Re \mathbf{X}} \right) \\
 &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) (\mathbf{A} \mathbf{X} (\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} + ((\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \mathbf{X}^H \mathbf{A})^T)
 \end{aligned}$$

Through the calculations, (16) and (35) were used. In addition, by use of (36), the derivative is found with respect to the imaginary part of \mathbf{X}

$$\begin{aligned}
 i \frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \Im \mathbf{X}} &= i \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) \left(\frac{\text{Tr}[\mathbf{A} \mathbf{X} (\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \partial(\mathbf{X}^H)]}{\partial \Im \mathbf{X}} + \frac{\text{Tr}[(\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \mathbf{X}^H \mathbf{A} \partial(\mathbf{X})]}{\partial \Im \mathbf{X}} \right) \\
 &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) (\mathbf{A} \mathbf{X} (\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} - ((\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \mathbf{X}^H \mathbf{A})^T)
 \end{aligned}$$

Hence, derivative yields

$$\begin{aligned} \frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \mathbf{X}} &= \frac{1}{2} \left(\frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \Re \mathbf{X}} - i \frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \Im \mathbf{X}} \right) \\ &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) ((\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \mathbf{X}^H \mathbf{A})^T \end{aligned}$$

and the complex conjugate derivative yields

$$\begin{aligned} \frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \mathbf{X}^*} &= \frac{1}{2} \left(\frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \Re \mathbf{X}} + i \frac{\partial \det(\mathbf{X}^H \mathbf{A} \mathbf{X})}{\partial \Im \mathbf{X}} \right) \\ &= \det(\mathbf{X}^H \mathbf{A} \mathbf{X}) \mathbf{A} \mathbf{X} (\mathbf{X}^H \mathbf{A} \mathbf{X})^{-1} \end{aligned}$$

Notice, for real \mathbf{X} , \mathbf{A} , the sum of (44) and (45) is reduced to (13).
Similar calculations yield

$$\begin{aligned} \frac{\partial \det(\mathbf{X} \mathbf{A} \mathbf{X}^H)}{\partial \mathbf{X}} &= \frac{1}{2} \left(\frac{\partial \det(\mathbf{X} \mathbf{A} \mathbf{X}^H)}{\partial \Re \mathbf{X}} - i \frac{\partial \det(\mathbf{X} \mathbf{A} \mathbf{X}^H)}{\partial \Im \mathbf{X}} \right) \\ &= \det(\mathbf{X} \mathbf{A} \mathbf{X}^H) (\mathbf{A} \mathbf{X}^H (\mathbf{X} \mathbf{A} \mathbf{X}^H)^{-1})^T \end{aligned} \quad (46)$$

and

$$\begin{aligned} \frac{\partial \det(\mathbf{X} \mathbf{A} \mathbf{X}^H)}{\partial \mathbf{X}^*} &= \frac{1}{2} \left(\frac{\partial \det(\mathbf{X} \mathbf{A} \mathbf{X}^H)}{\partial \Re \mathbf{X}} + i \frac{\partial \det(\mathbf{X} \mathbf{A} \mathbf{X}^H)}{\partial \Im \mathbf{X}} \right) \\ &= \det(\mathbf{X} \mathbf{A} \mathbf{X}^H) (\mathbf{X} \mathbf{A} \mathbf{X}^H)^{-1} \mathbf{X} \mathbf{A} \end{aligned} \quad (47)$$

References

- [1] Karl Gustav Andersson and Lars-Christer Boiers. *Ordinera differentialektioner*. Studentlitteratur, 1992.
- [2] Jörn Anemüller, Terrence J. Sejnowski, and Scott Makeig. Complex independent component analysis of frequency-domain electroencephalographic data. *Neural Networks*, 16(9):1311–1323, November 2003.
- [3] S. Barnett. *Matrices. Methods and Applications*. Oxford Applied Mathematics and Computing Science Series. Clarendon Press, 1990.
- [4] Christoffer Bishop. *Neural Networks for Pattern Recognition*. Oxford University Press, 1995.
- [5] Robert J. Boik. Lecture notes: Statistics 550. Online, April 22 2002. Notes.
- [6] D. H. Brandwood. A complex gradient operator and its application in adaptive array theory. *IEE Proceedings*, 130(1):11–16, February 1983. PTS. F and H.
- [7] M. Brookes. Matrix reference manual, 2004. Website May 20, 2004.
- [8] Mads Dyrholm. Some matrix results, 2004. Website August 23, 2004.
- [9] Simon Haykin. *Adaptive Filter Theory*. Prentice Hall, Upper Saddle River, NJ, 4th edition, 2002.
- [10] Thomas P. Minka. Old and new matrix algebra useful for statistics, December 2000. Notes.
- [11] L. Parra and C. Spence. Convolutional blind separation of non-stationary sources. In *IEEE Transactions Speech and Audio Processing*, pages 320–327, May 2000.
- [12] Laurent Schwartz. *Cours d'Analyse*, volume II. Hermann, Paris, 1967. As referenced in [9].
- [13] Shayle R. Searle. *Matrix Algebra Useful for Statistics*. John Wiley and Sons, 1982.
- [14] G. Seber and A. Lee. *Linear Regression Analysis*. John Wiley and Sons, 2002.
- [15] S. M. Selby. *Standard Mathematical Tables*. CRC Press, 1974.
- [16] Inna Stainvas. Matrix algebra in differential calculus. Neural Computing Research Group, Information Engineering, Aston University, UK, August 2002. Notes.
- [17] Max Welling. The kalman filter. Lecture Note.