


# Modeling Adult Visual Function

**Dr. James A. Bednar**

[jbednar@inf.ed.ac.uk](mailto:jbednar@inf.ed.ac.uk)

<http://homepages.inf.ed.ac.uk/jbednar>

# Surround modulation



Apparent contrast  
reduces


Detection facilitated

Contour pops out

Many types of contextual interactions are known

(Series et al. 2003)


# Surround modulation



Effects depend strongly on distance and contrast

Distance-related effects match both lateral and feedback connections

# Proposed model circuit




From Schwabe et al. (2006):

High-threshold inhibitory interneurons

Long-range excitatory lateral connections

Long-range excitatory feedback connections

# LESI circuit




From Law & Bednar (2006):

High-threshold inhibitory interneurons


Long-range excitatory lateral connections

No feedback connections yet

# Effective lateral inhibition



Excitatory activity



Inhibitory activity

(Law & Bednar 2006)


At high contrasts, the activity in the inhibitory sheet has wider radius than the activity in the excitatory sheet.

Result: Acts like Mexican-hat lateral interaction function, but using long-range excitatory connections.

Self-organization thus works as usual (since Hebbian learning is dominated by the high-contrast inputs), but circuitry is correct and low-contrast behavior can be correct.

# Stable development

## Standard LISSOM




(Law & Bednar 2006)

## Homeostatic no-shrinking laminar LISSOM

If the manual thresholds of standard LISSOM are replaced with homeostatic plasticity, excitatory radius shrinking can be eliminated. Result: map shape remains stable over time.


# The Tilt Aftereffect (TAE)



- Bias in orientation perception after prolonged exposure
- Allows model structure to be related to adult function




# TAE in Humans and LISSOM



- Mitchell & Muir 1976
- HLISSOM


- Direct effect for small angles
- Indirect effect for larger angles
- Model perception: vector average of orientations
- Human, model match closely

# TAE Adaptation in LISSOM




- **Adaptation:** More inhibition, but no net change in perception
- **Direct effect:** More inhibition for angles  $< 10^\circ$ 
  - Perception shifts from 10 to  $14^\circ$
- **Indirect effect:** Less inhibition for angles  $< 60^\circ$ 
  - Perception shifts from 60 to  $58^\circ$

# McCollough effect test pattern



Before adaptation,  
this pattern should  
appear monochrome

# Adaptation pattern




Stare alternately at the two patterns for 3 minutes,  
moving your gaze to avoid developing strong afterimages

# McCollough effect

(McCollough 1965)

After adaptation:




- Vertical bars should be slightly magenta
- Horizontal bars should be slightly green


- The effect should reverse if you tilt your head  $90^\circ$ , and disappear if you tilt  $45^\circ$ .

# McCollough effect: data

(Ellis 1977)




(Landisman & Ts'o 2002)



2.3 × 5.3mm macaque V1


- Effect measured in humans at each angle between adaptation and test
- Strength falls off smoothly with angle
- V1 is earliest possible substrate – first area showing OR selectivity; has color map

# LISSOM Color V1 Model



- Input: RGB images
- Decomposed into Red, Green channels (no blue in central fovea, Calkins 2001)
- Processed by color opponent retinal ganglia

# LISSOM OR + Color map



- Orientation map similar to animal maps
- Color-selective cells occur in blobs
- Each blob prefers either red or green




# Calculating McCollough Effect


- Perceived color estimated as a vector average of all units
- Vector direction: + for red-selective units, - for green-selective units
- Weighted by activation level and amount of color selectivity

Result is a number from extreme red (positive) to extreme green (negative), with approximately 0 being monochrome.

# Model McCollough Effect




# Compared with human



# Summary

- LISSOM can be compatible with actual circuit
- May explain surround modulation
- Afteffects arise from Hebbian adaptation of lateral inhibitory connections
- The same self-organizing processes can drive both development and adaptation: both structure and function
- **Novel prediction:** Indirect effect due to weight normalization

# McCullough Effect



Is the effect still  
present?

# References

- Bednar, J. A., De Paula, J. B., & Miikkulainen, R. (2005). Self-organization of color opponent receptive fields and laterally connected orientation maps. *Neurocomputing*, 65–66, 69–76.
- Calkins, D. J. (2001). Seeing with S cones. *Progress in Retinal and Eye Research*, 20 (3), 255–287.
- Ellis, S. R. (1977). Orientation selectivity of the McCollough effect: Analysis by equivalent contrast transformation. *Perception and Psychophysics*, 22 (6), 539–544.
- Landisman, C. E., & Ts'o, D. Y. (2002). Color processing in macaque striate cortex: Relationships to ocular dominance, cytochrome oxidase, and orientation. *Journal of Neurophysiology*, 87 (6), 3126–3137.

- Law, J. S., & Bednar, J. A. (2006). Surround modulation by long-range lateral connections in an orientation map model of primary visual cortex development and function. In *Society for Neuroscience Abstracts*. Society for Neuroscience, www.sfn.org. Program No. 546.4.
- McCollough, C. (1965). Color adaptation of edge-detectors in the human visual system. *Science*, *149* (3688), 1115–1116.
- Mitchell, D. E., & Muir, D. W. (1976). Does the tilt aftereffect occur in the oblique meridian?. *Vision Research*, *16*, 609–613.
- Schwabe, L., Obermayer, K., Angelucci, A., & Bressloff, P. C. (2006). The role of feedback in shaping the extra-classical receptive field of cortical neurons: A recurrent network model. *The Journal of Neuroscience*, *26* (36), 9117–9129.

Series, P., Lorenceau, J., & Fregnac, Y. (2003). The “silent” surround of V1 receptive fields: Theory and experiments. *Journal of Physiology (Paris)*, 97 (4–6), 453–474.