


Modeling Adult Visual Function

Dr. James A. Bednar

jbednar@inf.ed.ac.uk

http://homepages.inf.ed.ac.uk/jbednar

Surround modulation




(Series et al. 2003)

Apparent contrast reduces
Detection facilitated
Contour pops out

Many types of contextual interactions are known

Surround modulation




(Schwabe et al. 2006)

Effects depend strongly on distance and contrast

Distance-related effects match both lateral and feedback connections

Proposed model circuit



(Schwabe et al. 2006)


From Schwabe et al. (2006):

High-threshold inhibitory interneurons

Long-range excitatory lateral connections

Long-range excitatory feedback connections

LESI circuit




From Law & Bednar (2006):

High-threshold inhibitory interneurons


Long-range excitatory lateral connections

No feedback connections yet

Effective lateral inhibition



Excitatory activity



Inhibitory activity

(Law & Bednar 2006)


At high contrasts, the activity in the inhibitory sheet has wider radius than the activity in the excitatory sheet.

Result: Acts like Mexican-hat lateral interaction function, but using long-range excitatory connections.

Self-organization thus works as usual (since Hebbian learning is dominated by the high-contrast inputs), but circuitry is correct and low-contrast behavior can be correct.

Stable development

Standard LISSOM




(Law & Bednar 2006)

Homeostatic no-shrinking laminar LISSOM


If the manual thresholds of standard LISSOM are replaced with homeostatic plasticity, excitatory radius shrinking can be eliminated. Result: map shape remains stable over time.

The Tilt Aftereffect (TAE)



- Bias in orientation perception after prolonged exposure
- Allows model structure to be related to adult function


TAE in Humans and LISSOM



■ Mitchell & Muir 1976
■ HLISSOM


- Direct effect for small angles
- Indirect effect for larger angles
- Model perception: vector average of orientations
- Human, model match closely

TAE Adaptation in LISSOM




- **Adaptation:** More inhibition, but no net change in perception
- **Direct effect:** More inhibition for angles $< 10^\circ$
 - Perception shifts from 10 to 14°
- **Indirect effect:** Less inhibition for angles $< 60^\circ$
 - Perception shifts from 60 to 58°

McCullough effect test pattern




Before adaptation, this pattern should appear monochrome

Adaptation pattern



Stare alternately at the two patterns for 3 minutes, moving your gaze to avoid developing strong afterimages

McCollough effect





(McCollough 1965)

After adaptation:

- Vertical bars should be slightly magenta
- Horizontal bars should be slightly green

- The effect should reverse if you tilt your head 90°, and disappear if you tilt 45°.


McCollough effect: data



2.3x5.3mm macaque V1


- Effect measured in humans at each angle between adaptation and test
- Strength falls off smoothly with angle
- V1 is earliest possible substrate – first area showing OR selectivity; has color map

LISSOM Color V1 Model



- Input: RGB images
- Decomposed into Red, Green channels (no blue in central fovea, Calkins 2001)
- Processed by color opponent retinal ganglia

LISSOM OR + Color map




- Orientation map similar to animal maps
- Color-selective cells occur in blobs
- Each blob prefers either red or green

Calculating McCollough Effect


- Perceived color estimated as a vector average of all units
- Vector direction: + for red-selective units, - for green-selective units
- Weighted by activation level and amount of color selectivity

Result is a number from extreme red (positive) to extreme green (negative), with approximately 0 being monochrome.

Model McCollough Effect




Compared with human



Summary

- LISSOM can be compatible with actual circuit
- May explain surround modulation
- Aftffects arise from Hebbian adaptation of lateral inhibitory connections
- The same self-organizing processes can drive both development and adaptation: both structure and function
- **Novel prediction:** Indirect effect due to weight normalization

McCollough Effect



Is the effect still present?

Law, J. S., & Bednar, J. A. (2006). Surround modulation by long-range lateral connections in an orientation map model of primary visual cortex development and function. In *Society for Neuroscience Abstracts*. Society for Neuroscience, www.sfn.org. Program No. 546.4.

McCollough, C. (1965). Color adaptation of edge-detectors in the human visual system. *Science*, 149 (3688), 1115–1116.

Mitchell, D. E., & Muir, D. W. (1976). Does the tilt aftereffect occur in the oblique meridian?. *Vision Research*, 16, 609–613.

Schwabe, L., Obermayer, K., Angelucci, A., & Bressloff, P. C. (2006). The role of feedback in shaping the extra-classical receptive field of cortical neurons: A recurrent network model. *The Journal of Neuroscience*, 26 (36), 9117–9129.

References

- Bednar, J. A., De Paula, J. B., & Miikkulainen, R. (2005). Self-organization of color opponent receptive fields and laterally connected orientation maps. *Neurocomputing*, 65–66, 69–76.
- Calkins, D. J. (2001). Seeing with S cones. *Progress in Retinal and Eye Research*, 20 (3), 255–287.
- Ellis, S. R. (1977). Orientation selectivity of the McCollough effect: Analysis by equivalent contrast transformation. *Perception and Psychophysics*, 22 (6), 539–544.
- Landisman, C. E., & Ts'o, D. Y. (2002). Color processing in macaque striate cortex: Relationships to ocular dominance, cytochrome oxidase, and orientation. *Journal of Neurophysiology*, 87 (6), 3126–3137.

Series, P., Lorenceau, J., & Fregnac, Y. (2003). The “silent” surround of V1 receptive fields: Theory and experiments. *Journal of Physiology (Paris)*, 97 (4–6), 453–474.