


# Modeling Adult Visual Function

**Dr. James A. Bednar**

[jbednar@inf.ed.ac.uk](mailto:jbednar@inf.ed.ac.uk)

<http://homepages.inf.ed.ac.uk/jbednar>

# Surround modulation



Apparent contrast  
reduces

Detection facilitated


Contour pops out

Many types of contextual interactions are known

(Series et al. 2003)

# Surround modulation

(Schwabe et al. 2006)



Effects depend strongly on contrast (Hirsch & Gilbert 1991), (Weliky et al. 1995) and on distance

Distance-related effects match both lateral and feedback connections

Distance-related effects match both lateral and feedback connections


Distance-related effects match both lateral and feedback connections

Distance-related effects match both lateral and feedback connections

Distance-related effects match both lateral and feedback connections

Distance-related effects match both lateral and feedback connections

# Proposed model circuit




From Schwabe et al. (2006):

High-threshold inhibitory interneurons

Long-range excitatory lateral connections

Long-range excitatory feedback connections

# GCAL SM model




(Antolik 2010; Antolik & Bednar 2012)

- GCAL circuit for surround modulation
- Separate inhibitory interneurons
- Long-range excitatory lateral connections
- Separate simple and complex cell layers
- Feedback connections in progress (Philipp Rudiger)


# SM model size tuning

(Antolik 2010)




Single-unit response to larger patterns typically increases,  
then decreases as inhibition is recruited

# Diversity in size tuning



Model matches both typical and unusual size tuning responses


# Diversity in OCTC tuning



Model matches both typical and unusual orientation-contrast tuning types




# The Tilt Aftereffect (TAE)




- Bias in orientation perception after prolonged exposure
- Allows model structure to be related to adult function
- Classic explanation: “fatigue” – activated neurons get tired, shifting the population average away

# Measuring perceived orientation



- Assumption: perception based on population average
- Vector average good for cyclic quantities
- Decode perception before and after adaptation


# TAE in Humans and LISSOM



■ Mitchell & Muir 1976  
■ HLISSOM


- Direct effect for small angles
- Indirect effect for larger angles
- Null effect at training angle
- Human, model match closely

# TAE Adaptation in LISSOM




- **Null at zero:** More inhibition, but no net change in perception
- **Direct effect:** More inhibition for angles  $< 10^\circ$ 
  - Perception shifts from 10 to 14°
- **Indirect effect:** Less inhibition for angles  $< 60^\circ$ 
  - Perception shifts from 60 to 58°
- Due to synapses, not tired neurons!

# McCollough effect test pattern



Before adaptation,  
this pattern should  
appear monochrome

# Adaptation pattern




Stare alternately at the two patterns for 3 minutes,  
moving your gaze to avoid developing strong afterimages

# McCollough effect

(McCollough 1965)

After adaptation:




- Vertical bars should be slightly magenta
- Horizontal bars should be slightly green


- The effect should reverse if you tilt your head  $90^\circ$ , and disappear if you tilt  $45^\circ$ .

# McCollough effect: data

(Ellis 1977)



(Landisman & Ts'o 2002)




2.3 × 5.3mm macaque V1

- Effect measured in humans at each angle between adaptation and test
- Strength falls off smoothly with angle
- V1 is earliest possible substrate – first area showing OR selectivity; has color map




# LISSOM Color V1 Model



- Input: RGB images
- Decomposed into Red, Green channels (no blue in central fovea, [Calkins 2001](#))
- Processed by color opponent retinal ganglia

# LISSOM OR + Color map




- Orientation map similar to animal maps
- Color-selective cells occur in blobs
- Preferences of neurons in each blob?

# Calculating McCollough Effect


- Perceived color estimated as a vector average of all units
- Vector direction: + for red-selective units, - for green-selective units
- Weighted by activation level and amount of color selectivity

Result is a number from extreme red (positive) to extreme green (negative), with approximately 0 being monochrome.

# Model McCollough Effect




# Compared with human



# Summary

- GCAL can be compatible with actual circuit
- Reproduces surprising features of surround modulation
- Afteffects arise from Hebbian adaptation of lateral connections
- The same self-organizing processes can drive both development and adaptation: both structure and function
- **Novel prediction:** Indirect effect due to weight normalization
- Project: exactly how does inverted Mexican Hat work?

# McCullough Effect



Is the effect still  
present?

# References

- Antolik, J. (2010). *Unified Developmental Model of Maps, Complex Cells and Surround Modulation in the Primary Visual Cortex*. Doctoral Dissertation, School of Informatics, The University of Edinburgh, Edinburgh, UK.
- Antolik, J., & Bednar, J. A. (2012). A unified developmental model of maps, complex cells and surround modulation in the primary visual cortex. In preparation.
- Bednar, J. A., De Paula, J. B., & Miikkulainen, R. (2005). Self-organization of color opponent receptive fields and laterally connected orientation maps. *Neurocomputing*, 65–66, 69–76.
- Calkins, D. J. (2001). Seeing with S cones. *Progress in Retinal and Eye Research*, 20 (3), 255–287.



- Ellis, S. R. (1977). Orientation selectivity of the McCollough effect: Analysis by equivalent contrast transformation. *Perception and Psychophysics*, *22* (6), 539–544.
- Hirsch, J. A., & Gilbert, C. D. (1991). Synaptic physiology of horizontal connections in the cat's visual cortex. *The Journal of Neuroscience*, *11*, 1800–1809.
- Landisman, C. E., & Ts'o, D. Y. (2002). Color processing in macaque striate cortex: Relationships to ocular dominance, cytochrome oxidase, and orientation. *Journal of Neurophysiology*, *87* (6), 3126–3137.
- McCollough, C. (1965). Color adaptation of edge-detectors in the human visual system. *Science*, *149* (3688), 1115–1116.
- Mitchell, D. E., & Muir, D. W. (1976). Does the tilt aftereffect occur in the oblique meridian?. *Vision Research*, *16*, 609–613.

- Schwabe, L., Obermayer, K., Angelucci, A., & Bressloff, P. C. (2006). The role of feedback in shaping the extra-classical receptive field of cortical neurons: A recurrent network model. *The Journal of Neuroscience*, *26* (36), 9117–9129.
- Series, P., Lorenceau, J., & Fregnac, Y. (2003). The “silent” surround of V1 receptive fields: Theory and experiments. *Journal of Physiology (Paris)*, *97* (4–6), 453–474.
- Weliky, M., Kandler, K., Fitzpatrick, D., & Katz, L. C. (1995). Patterns of excitation and inhibition evoked by horizontal connections in visual cortex share a common relationship to orientation columns. *Neuron*, *15*, 541–552.