


Exercises for the tutorials: 1, 2(a-b), 3.

The other exercises are for self-study and exam preparation. All material is examinable unless otherwise mentioned.

**Exercise 1. Directed graph concepts**

We here consider the directed graph below that was partly discussed in the lecture.



(a) List all trails in the graph (of maximal length)

**Solution.** We have

$$(a, q, e) \quad (a, q, z, h) \quad (h, z, q, e)$$

and the corresponding ones with swapped start and end nodes.

(b) List all directed paths in the graph (of maximal length)

**Solution.**  $(a, q, e)$      $(z, q, e)$      $(z, h)$

(c) What are the descendants of  $z$ ?

**Solution.**  $\text{desc}(z) = \{q, e, h\}$

(d) What are the non-descendants of  $q$ ?

**Solution.**  $\text{nondesc}(q) = \{a, z, h, e\} \setminus \{e\} = \{a, z, h\}$

(e) Which of the following orderings are topological to the graph?

- $(a, z, h, q, e)$
- $(a, z, e, h, q)$
- $(z, a, q, h, e)$
- $(z, q, e, a, h)$

**Solution.**

- $(a, z, h, q, e)$ : yes
- $(a, z, e, h, q)$ : no ( $q$  is a parent of  $e$  and thus has to come before  $e$  in the ordering)
- $(z, a, q, h, e)$ : yes
- $(z, q, e, a, h)$ : no ( $a$  is a parent of  $q$  and thus has to come before  $q$  in the ordering)

## Exercise 2. Canonical connections

We here derive the independencies that hold in the three canonical connections that exist in DAGs, shown in Figure 1.




Figure 1: The three canonical connections in DAGs.

(a) For the serial connection, use the ordered Markov property to show that  $x \perp\!\!\!\perp y \mid z$ .

**Solution.** The only topological ordering is  $x, z, y$ . The predecessors of  $y$  are  $\text{pre}_y = \{x, z\}$  and its parents  $\text{pa}_y = \{z\}$ . The ordered Markov property

$$y \perp\!\!\!\perp (\text{pre}_y \setminus \text{pa}_y) \mid \text{pa}_y \tag{S.1}$$

thus becomes  $y \perp\!\!\!\perp (\{x, z\} \setminus z) \mid z$ . Hence we have

$$y \perp\!\!\!\perp x \mid z, \tag{S.2}$$

which is the same as  $x \perp\!\!\!\perp y \mid z$  since the independency relationship is symmetric.

This means that if the state or value of  $z$  is known (i.e. if the random variable  $z$  is “instantiated”), evidence about  $x$  will not change our belief about  $y$ , and vice versa. We say that the  $z$  node is “closed” and that the trail between  $x$  and  $y$  is “blocked” by the instantiated  $z$ . In other words, knowing the value of  $z$  blocks the flow of evidence *between*  $x$  and  $y$ .

(b) For the serial connection, show that the marginal  $p(x, y)$  does generally not factorise into  $p(x)p(y)$ , i.e. that  $x \perp\!\!\!\perp y$  does not hold.

**Solution.** There are several ways to show the result. One is to present an example where the independency does not hold. Consider for instance the following model

$$x \sim \mathcal{N}(x; 0, 1) \tag{S.3}$$

$$z = x + n_z \tag{S.4}$$

$$y = z + n_y \tag{S.5}$$

where  $n_z \sim \mathcal{N}(n_z; 0, 1)$  and  $n_y \sim \mathcal{N}(n_y; 0, 1)$ , both being statistically independent from  $x$ . Here  $\mathcal{N}(\cdot; 0, 1)$  denotes the Gaussian pdf with mean 0 and variance 1, and  $x \sim \mathcal{N}(x; 0, 1)$  means that we sample  $x$  from the distribution  $\mathcal{N}(x; 0, 1)$ . Hence  $p(z|x) = \mathcal{N}(z; x, 1)$ ,  $p(y|z) = \mathcal{N}(y; z, 1)$  and  $p(x, y, z) = p(x)p(z|x)p(y|z) = \mathcal{N}(x; 0, 1)\mathcal{N}(z; x, 1)\mathcal{N}(y; z, 1)$ .

Whilst we could manipulate the pdfs to show the result, it’s here easier to work with the generative model in Equations (S.3) to (S.5). Eliminating  $z$  from the equations, by plugging the definition of  $z$  into (S.5) we have

$$y = x + n_z + n_y, \tag{S.6}$$

which describes the marginal distribution of  $(x, y)$ . We see that  $\mathbb{E}[xy]$  is

$$\mathbb{E}[xy] = \mathbb{E}[x^2 + xn_z + xn_y] \quad (\text{S.7})$$

$$= \mathbb{E}[x^2] + \mathbb{E}[x]\mathbb{E}[n_z] + \mathbb{E}[x]\mathbb{E}[n_y] \quad (\text{S.8})$$

$$= 1 + 0 + 0 \quad (\text{S.9})$$

where we have used the linearity of expectation, that  $x$  is independent from  $n_z$  and  $n_y$ , and that  $x$  has zero mean. If  $x$  and  $y$  were independent (or only uncorrelated), we had  $\mathbb{E}[xy] = \mathbb{E}[x]\mathbb{E}[y] = 0$ . However, since  $\mathbb{E}[xy] \neq \mathbb{E}[x]\mathbb{E}[y]$ ,  $x$  and  $y$  are not independent.

In plain English, this means that if the state of  $z$  is unknown, then evidence or information about  $x$  will influence our belief about  $y$ , and the other way around. Evidence can flow through  $z$  between  $x$  and  $y$ . We say that the  $z$  node is “open” and the trail between  $x$  and  $y$  is “active”.

(c) For the diverging connection, use the ordered Markov property to show that  $x \perp\!\!\!\perp y \mid z$ .

**Solution.** A topological ordering is  $z, x, y$ . The predecessors of  $y$  are  $\text{pre}_y = \{x, z\}$  and its parents  $\text{pa}_y = \{z\}$ . The ordered Markov property

$$y \perp\!\!\!\perp (\text{pre}_y \setminus \text{pa}_y) \mid \text{pa}_y \quad (\text{S.10})$$

thus becomes again

$$y \perp\!\!\!\perp x \mid z, \quad (\text{S.11})$$

which is, since the independence relationship is symmetric, the same as  $x \perp\!\!\!\perp z \mid y$ .

As in the serial connection, if the state or value  $z$  is known, evidence about  $x$  will not change our belief about  $y$ , and vice versa. Knowing  $z$  closes the  $z$  node, which blocks the trail between  $x$  and  $y$ .

(d) For the diverging connection, show that the marginal  $p(x, y)$  does generally not factorise into  $p(x)p(y)$ , i.e. that  $x \perp\!\!\!\perp y$  does not hold.

**Solution.** As for the serial connection, it suffices to give an example where  $x \perp\!\!\!\perp y$  does not hold. We consider the following generative model

$$z \sim \mathcal{N}(z; 0, 1) \quad (\text{S.12})$$

$$x = z + n_x \quad (\text{S.13})$$

$$y = z + n_y \quad (\text{S.14})$$

where  $n_x \sim \mathcal{N}(n_x; 0, 1)$  and  $n_y \sim \mathcal{N}(n_y; 0, 1)$ , and they are independent of each other and the other variables. We have  $\mathbb{E}[x] = \mathbb{E}[z + n_x] = \mathbb{E}[z] + \mathbb{E}[n_x] = 0$ . On the other hand

$$\mathbb{E}[xy] = \mathbb{E}[(z + n_x)(z + n_y)] \quad (\text{S.15})$$

$$= \mathbb{E}[z^2 + z(n_x + n_y) + n_x n_y] \quad (\text{S.16})$$

$$= \mathbb{E}[z^2] + \mathbb{E}[z(n_x + n_y)] + \mathbb{E}[n_x n_y] \quad (\text{S.17})$$

$$= 1 + 0 + 0 \quad (\text{S.18})$$

Hence,  $\mathbb{E}[xy] \neq \mathbb{E}[x]\mathbb{E}[y]$  and we do not have that  $x \perp\!\!\!\perp y$  holds.

In a diverging connection, as in the serial connection, if the state of  $z$  is unknown, then evidence or information about  $x$  will influence our belief about  $y$ , and the other way around. Evidence can flow through  $z$  between  $x$  and  $y$ . We say that the  $z$  node is open and the trail between  $x$  and  $y$  is active.

(e) For the converging connection, show that  $x \perp\!\!\!\perp y$ .

**Solution.** We can here again use the ordered Markov property with the ordering  $y, x, z$ . Since  $\text{pre}_x = \{y\}$  and  $\text{pa}_x = \emptyset$ , we have

$$x \perp\!\!\!\perp (\text{pre}_x \setminus \text{pa}_x) \mid \text{pa}_x = x \perp\!\!\!\perp y. \quad (\text{S.19})$$

Alternatively, we can use the basic definition of directed graphical models, i.e.

$$p(x, y, z) = k(x)k(y)k(z \mid x, y) \quad (\text{S.20})$$

together with the result that the kernels (factors) are valid (conditional) pdfs/pmfs and equal to the conditionals/marginals with respect to the joint distribution  $p(x, y, z)$ , i.e.

$$k(x) = p(x) \quad (\text{S.21})$$

$$k(y) = p(y) \quad (\text{S.22})$$

$$k(z \mid x, y) = p(z \mid x, y) \quad (\text{not needed in the proof below}) \quad (\text{S.23})$$

Integrating out  $z$  gives

$$p(x, y) = \int p(x, y, z) dz \quad (\text{S.24})$$

$$= \int k(x)k(y)k(z \mid x, y) dz \quad (\text{S.25})$$

$$= k(x)k(y) \underbrace{\int k(z \mid x, y) dz}_1 \quad (\text{S.26})$$

$$= p(x)p(y) \quad (\text{S.27})$$

Hence  $p(x, y)$  factorises into its marginals, which means that  $x \perp\!\!\!\perp y$ .

Hence, when we do not have evidence about  $z$ , evidence about  $x$  will not change our belief about  $y$ , and vice versa. For the converging connection, if no evidence about  $z$  is available, the  $z$  node is closed, which blocks the trail between  $x$  and  $y$ .

(f) For the converging connection, show that  $x \perp\!\!\!\perp y \mid z$  does generally not hold.

**Solution.** We give a simple example where  $x \perp\!\!\!\perp y \mid z$  does not hold.

Consider

$$x \sim \mathcal{N}(x; 0, 1) \quad (\text{S.28})$$

$$y \sim \mathcal{N}(y; 0, 1) \quad (\text{S.29})$$

$$z = xy + n_z \quad (\text{S.30})$$

where  $n_z \sim \mathcal{N}(n_z; 0, 1)$ , independent from the other variables. From the last equation, we have

$$xy = z - n_z \quad (\text{S.31})$$

We thus have

$$\mathbb{E}[xy \mid z] = \mathbb{E}[z - n_z \mid z] \quad (\text{S.32})$$

$$= z - 0 \quad (\text{S.33})$$

On the other hand,  $\mathbb{E}[xy] = \mathbb{E}[x]\mathbb{E}[y] = 0$ . Since  $\mathbb{E}[xy \mid z] \neq \mathbb{E}[xy]$ ,  $x \perp\!\!\!\perp y \mid z$  cannot hold.


The intuition here is that if you know the value of the product  $xy$ , even if subject to noise, knowing the value of  $x$  allows you to guess the value of  $y$  and vice versa.

More generally, for converging connections, if evidence or information about  $z$  is available, evidence about  $x$  will influence the belief about  $y$ , and vice versa. We say that information about  $z$  opens the  $z$ -node, and evidence can flow between  $x$  and  $y$ .

Note: information about  $z$  means that  $z$  or one of its descendants is observed, see exercise 9.

### Exercise 3. Ordered and local Markov properties, d-separation

We continue with the investigation of the graph from Exercise 1 shown below for reference.



- (a) The ordering  $(z, h, a, q, e)$  is topological to the graph. What are the independencies that follow from the ordered Markov property?

**Solution.** We proceed as in the lecture slides: The predecessor sets are

$$\text{pre}_z = \emptyset, \text{pre}_h = \{z\}, \text{pre}_a = \{z, h\}, \text{pre}_q = \{z, h, a\}, \text{pre}_e = \{z, h, a, q\}$$

The parent sets are independent from the topological ordering chosen. In the lecture, we have seen that they are:

$$\text{pa}_z = \emptyset, \text{pa}_h = \{z\}, \text{pa}_a = \emptyset, \text{pa}_q = \{a, z\}, \text{pa}_e = \{q\},$$

The ordered Markov property reads  $x_i \perp\!\!\!\perp (\text{pre}_i \setminus \text{pa}_i) \mid \text{pa}_i$  where the  $x_i$  refer to the ordered variables, e.g.  $x_1 = z, x_2 = h, x_3 = a$ , etc.

With

$$\text{pre}_h \setminus \text{pa}_h = \emptyset \quad \text{pre}_a \setminus \text{pa}_a = \{z, h\} \quad \text{pre}_q \setminus \text{pa}_q = \{h\} \quad \text{pre}_e \setminus \text{pa}_e = \{z, h, a\}$$

we thus obtain

$$h \perp\!\!\!\perp \emptyset \mid z \quad a \perp\!\!\!\perp \{z, h\} \quad q \perp\!\!\!\perp h \mid \{a, z\} \quad e \perp\!\!\!\perp \{z, h, a\} \mid q$$

The relation  $h \perp\!\!\!\perp \emptyset \mid z$  should be understood as “there is no variable from which  $h$  is independent given  $z$ ” and should thus be dropped from the list. Compared to the relations obtained for the orderings in the lecture, the new one here is  $a \perp\!\!\!\perp \{z, h\}$ . Generally, having a variable later in the topological ordering allows one to possibly obtain a stronger independence relation because the set  $\text{pre} \setminus \text{pa}$  can only increase when the predecessor set  $\text{pre}$  becomes larger.

- (b) What are the independencies that follow from the local Markov property?

**Solution.** The non-descendants are

$$\text{nondesc}(a) = \{z, h\} \quad \text{nondesc}(z) = \{a\} \quad \text{nondesc}(h) = \{a, z, q, e\}$$

$$\text{nondesc}(q) = \{a, z, h\} \quad \text{nondesc}(e) = \{a, q, z, h\}$$

With the parent sets as before, the independencies that follow from the local Markov property are  $x_i \perp\!\!\!\perp (\text{nondesc}(x_i) \setminus \text{pa}_i) \mid \text{pa}_i$ , i.e.


$$a \perp\!\!\!\perp \{z, h\} \quad z \perp\!\!\!\perp a \quad h \perp\!\!\!\perp \{a, q, e\} \mid z \quad q \perp\!\!\!\perp h \mid \{a, z\} \quad e \perp\!\!\!\perp \{a, z, h\} \mid q$$

- (c) The independency relations obtained via the ordered and local Markov property include  $q \perp\!\!\!\perp h \mid \{a, z\}$ . Verify the independency using d-separation.


**Solution.** The only trail from  $q$  to  $h$  goes through  $z$  which is in a tail-tail configuration. Since  $z$  is part of the conditioning set, the trail is blocked and the result follows.

- (d) Use d-separation to check whether  $a \perp\!\!\!\perp h \mid e$  holds.

**Solution.** The trail from  $a$  to  $h$  is shown below in red together with the default states of the nodes along the trail.



Conditioning on  $e$  opens the  $q$  node since  $q$  is in a collider configuration on the path.



The trail from  $a$  to  $h$  is thus active, which means that the relationship does not hold because  $a \not\perp\!\!\!\perp h \mid e$  for some distributions that factorise over the graph.

- (e) Assume all variables in the graph are binary. How many numbers do you need to specify, or learn from data, in order to fully specify the probability distribution?

**Solution.** The graph defines a set of probability mass functions (pmf) that factorise as

$$p(a, z, q, h, e) = p(a)p(z)p(q|a, z)p(h|z)p(e|q)$$


To specify a member of the set, we need to specify the (conditional) pmfs on the right-hand side. The (conditional) pmfs can be seen as tables, and the number of elements that we need to specified in the tables are:

- 1 for  $p(a)$
- 1 for  $p(z)$
- 4 for  $p(q|a, z)$
- 2 for  $p(h|z)$
- 2 for  $p(e|q)$

In total, there are 10 numbers to specify. This is in contrast to  $2^5 - 1 = 31$  for a distribution without independencies. Note that the number of parameters to specify could be further reduced by making parametric assumptions.

**Exercise 4. More on ordered and local Markov properties, d-separation**

We continue with the investigation of the graph below



- (a) Why can the ordered or local Markov property not be used to check whether  $a \perp\!\!\!\perp h \mid e$  may hold?

**Solution.** The independencies that follow from the ordered or local Markov property require conditioning on parent sets. However,  $e$  is not a parent of any node so that the above independence assertion cannot be checked via the ordered or local Markov property.

- (b) The independency relations obtained via the ordered and local Markov property include  $a \perp\!\!\!\perp \{z, h\}$ . Verify the independency using d-separation.

**Solution.** All paths from  $a$  to  $z$  or  $h$  pass through the node  $q$  that forms a head-head connection along that trail. Since neither  $q$  nor its descendant  $e$  is part of the conditioning set, the trail is blocked and the independence relation follows.

- (c) Determine the Markov blanket of  $z$ .

**Solution.** The Markov blanket is given by the parents, children, and co-parents. Hence:  $MB(z) = \{a, q, h\}$ .

- (d) Verify that  $q \perp\!\!\!\perp h \mid \{a, z\}$  holds by manipulating the probability distribution induced by the graph.

**Solution.** A basic definition of conditional statistical independence  $x_1 \perp\!\!\!\perp x_2 \mid x_3$  is that the (conditional) joint  $p(x_1, x_2 \mid x_3)$  equals the product of the (conditional) marginals  $p(x_1 \mid x_3)$  and  $p(x_2 \mid x_3)$ . In other words, for discrete random variables,

$$x_1 \perp\!\!\!\perp x_2 \mid x_3 \iff p(x_1, x_2 \mid x_3) = \left( \sum_{x_2} p(x_1, x_2 \mid x_3) \right) \left( \sum_{x_1} p(x_1, x_2 \mid x_3) \right) \quad (\text{S.34})$$

We thus answer the question by showing that (use integrals in case of continuous random variables)

$$p(q, h \mid a, z) = \left( \sum_h p(q, h \mid a, z) \right) \left( \sum_q p(q, h \mid a, z) \right) \quad (\text{S.35})$$

First, note that the graph defines a set of probability density or mass functions that factorise as

$$p(a, z, q, h, e) = p(a)p(z)p(q \mid a, z)p(h \mid z)p(e \mid q)$$

We then use the sum-rule to compute the joint distribution of  $(a, z, q, h)$ , i.e. the distribution of all the variables that occur in  $p(q, h \mid a, z)$

$$p(a, z, q, h) = \sum_e p(a, z, q, h, e) \quad (\text{S.36})$$

$$= \sum_e p(a)p(z)p(q \mid a, z)p(h \mid z)p(e \mid q) \quad (\text{S.37})$$

$$= p(a)p(z)p(q \mid a, z)p(h \mid z) \underbrace{\sum_e p(e \mid q)}_1 \quad (\text{S.38})$$

$$= p(a)p(z)p(q \mid a, z)p(h \mid z), \quad (\text{S.39})$$

where  $\sum_e p(e \mid q) = 1$  because (conditional) pdfs/pmfs are normalised so that the integrate/sum to one. We further have

$$p(a, z) = \sum_{q, h} p(a, z, q, h) \quad (\text{S.40})$$

$$= \sum_{q, h} p(a)p(z)p(q \mid a, z)p(h \mid z) \quad (\text{S.41})$$

$$= p(a)p(z) \sum_q p(q \mid a, z) \sum_h p(h \mid z) \quad (\text{S.42})$$

$$= p(a)p(z) \quad (\text{S.43})$$

so that

$$p(q, h \mid a, z) = \frac{p(a, z, q, h)}{p(a, z)} \quad (\text{S.44})$$

$$= \frac{p(a)p(z)p(q \mid a, z)p(h \mid z)}{p(a)p(z)} \quad (\text{S.45})$$

$$= p(q \mid a, z)p(h \mid z). \quad (\text{S.46})$$

We further see that  $p(q \mid a, z)$  and  $p(h \mid z)$  are the marginals of  $p(q, h \mid a, z)$ , i.e.

$$p(q \mid a, z) = \sum_h p(q, h \mid a, z) \quad (\text{S.47})$$

$$p(h \mid z) = \sum_q p(q, h \mid a, z). \quad (\text{S.48})$$



This means that

$$p(q, h|a, z) = \left( \sum_h p(q, h|a, z) \right) \left( \sum_q p(q, h|a, z) \right), \quad (\text{S.49})$$

which shows that  $q \perp\!\!\!\perp h|a, z$ .

We see that using the graph to determine the independency is easier than manipulating the pmf/pdf.

**Exercise 5. Chest clinic (based on Barber’s exercise 3.3)**

The directed graphical model in Figure 2 is about the diagnosis of lung disease ( $t$ =tuberculosis or  $l$ =lung cancer). In this model, a visit to some place “ $a$ ” is thought to increase the probability of tuberculosis.




Figure 2: Graphical model for Exercise 5 (Barber Figure 3.15).

(a) Explain which of the following independence relationships hold for all distributions that factorise over the graph.

1.  $t \perp\!\!\!\perp s \mid d$

**Solution.**

- There are two trails from  $t$  to  $s$ :  $(t, e, l, s)$  and  $(t, e, d, b, s)$ .
- The trail  $(t, e, l, s)$  features a collider node  $e$  that is opened by the conditioning variable  $d$ . The trail is thus active and we do not need to check the second trail because for independence all trails needed to be blocked.
- The independence relationship does thus generally not hold.

2.  $l \perp\!\!\!\perp b \mid s$

**Solution.**

- There are two trails from  $l$  to  $b$ :  $(l, s, b)$  and  $(l, e, d, b)$
- The trail  $(l, s, b)$  is blocked by  $s$  ( $s$  is in a tail-tail configuration and part of the conditioning set)
- The trail  $(l, e, d, b)$  is blocked by the collider configuration for node  $d$ .
- All trails are blocked so that the independence relation holds.

(b) Can we simplify  $p(l|b, s)$  to  $p(l|s)$ ?

**Solution.** Since  $l \perp\!\!\!\perp b \mid s$ , we have  $p(l|b, s) = p(l|s)$ .

**Exercise 6. More on the chest clinic (based on Barber's exercise 3.3)**

Consider the directed graphical model in Figure 2.

(a) Explain which of the following independence relationships hold for all distributions that factorise over the graph.

1.  $a \perp\!\!\!\perp s \mid l$

**Solution.**

- There are two trails from  $a$  to  $s$ :  $(a, t, e, l, s)$  and  $(a, t, e, d, b, s)$
- The trail  $(a, t, e, l, s)$  features a collider node  $e$  that blocks the trail (the trail is also blocked by  $l$ ).
- The trail  $(a, t, e, d, b, s)$  is blocked by the collider node  $d$ .
- All trails are blocked so that the independence relation holds.

2.  $a \perp\!\!\!\perp s \mid l, d$

**Solution.**

- There are two trails from  $a$  to  $s$ :  $(a, t, e, l, s)$  and  $(a, t, e, d, b, s)$
- The trail  $(a, t, e, l, s)$  features a collider node  $e$  that is opened by the conditioning variable  $d$  but the  $l$  node is closed by the conditioning variable  $l$ : the trail is blocked
- The trail  $(a, t, e, d, b, s)$  features a collider node  $d$  that is opened by conditioning on  $d$ . On this trail,  $e$  is not in a head-head (collider) configuration) so that all nodes are open and the trail active.
- Hence, the independence relation does generally not hold.

(b) Let  $g$  be a (deterministic) function of  $x$  and  $t$ . Is the expected value  $\mathbb{E}[g(x, t) \mid l, b]$  equal to  $\mathbb{E}[g(x, t) \mid l]$ ?


**Solution.** The question boils down to checking whether  $x, t \perp\!\!\!\perp b \mid l$ . For the independence relation to hold, all trails from both  $x$  and  $t$  to  $b$  need to be blocked by  $l$ .

- For  $x$ , we have the trails  $(x, e, l, s, b)$  and  $(x, e, d, b)$
- Trail  $(x, e, l, s, b)$  is blocked by  $l$
- Trail  $(x, e, d, b)$  is blocked by the collider configuration of node  $d$ .
- For  $t$ , we have the trails  $(t, e, l, s, b)$  and  $(t, e, d, b)$
- Trail  $(t, e, l, s, b)$  is blocked by  $l$ .
- Trail  $(t, e, d, b)$  is blocked by the collider configuration of node  $d$ .

As all trails are blocked we have  $x, t \perp\!\!\!\perp b \mid l$  and  $\mathbb{E}[g(x, t) \mid l, b] = \mathbb{E}[g(x, t) \mid l]$ .

### Exercise 7. Hidden Markov models

This exercise is about directed graphical models that are specified by the following DAG:



These models are called “hidden” Markov models because we typically assume to only observe the  $y_i$  and not the  $x_i$  that follow a Markov model.

(a) Show that all probabilistic models specified by the DAG factorise as

$$p(x_1, y_1, x_2, y_2, \dots, x_4, y_4) = p(x_1)p(y_1|x_1)p(x_2|x_1)p(y_2|x_2)p(x_3|x_2)p(y_3|x_3)p(x_4|x_3)p(y_4|x_4)$$

**Solution.** From the definition of directed graphical models it follows that

$$p(x_1, y_1, x_2, y_2, \dots, x_4, y_4) = \prod_{i=1}^4 p(x_i | \text{pa}(x_i)) \prod_{i=1}^4 p(y_i | \text{pa}(y_i)).$$

The result is then obtained by noting that the parent of  $y_i$  is given by  $x_i$  for all  $i$ , and that the parent of  $x_i$  is  $x_{i-1}$  for  $i = 2, 3, 4$  and that  $x_1$  does not have a parent ( $\text{pa}(x_1) = \emptyset$ ).

(b) Derive the independencies implied by the ordered Markov property with the topological ordering  $(x_1, y_1, x_2, y_2, x_3, y_3, x_4, y_4)$

**Solution.**

$$y_i \perp\!\!\!\perp x_1, y_1, \dots, x_{i-1}, y_{i-1} \mid x_i \quad x_i \perp\!\!\!\perp x_1, y_1, \dots, x_{i-2}, y_{i-2}, y_{i-1} \mid x_{i-1}$$

(c) Derive the independencies implied by the ordered Markov property with the topological ordering  $(x_1, x_2, \dots, x_4, y_1, \dots, y_4)$ .

**Solution.** For the  $x_i$ , we use that for  $i \geq 2$ :  $\text{pre}(x_i) = \{x_1, \dots, x_{i-1}\}$  and  $\text{pa}(x_i) = x_{i-1}$ . For the  $y_i$ , we use that  $\text{pre}(y_1) = \{x_1, \dots, x_4\}$ , that  $\text{pre}(y_i) = \{x_1, \dots, x_4, y_1, \dots, y_{i-1}\}$  for  $i > 1$ , and that  $\text{pa}(y_i) = x_i$ . The ordered Markov property then gives:

$$\begin{array}{ll} x_3 \perp\!\!\!\perp x_1 \mid x_2 & x_4 \perp\!\!\!\perp \{x_1, x_2\} \mid x_3 \\ y_1 \perp\!\!\!\perp \{x_2, x_3, x_4\} \mid x_1 & y_2 \perp\!\!\!\perp \{x_1, x_3, x_4, y_1\} \mid x_2 \\ y_3 \perp\!\!\!\perp \{x_1, x_2, x_4, y_1, y_2\} \mid x_3 & y_4 \perp\!\!\!\perp \{x_1, x_2, x_3, y_1, y_2, y_3\} \mid x_4 \end{array}$$

(d) Does  $y_4 \perp\!\!\!\perp y_1 \mid y_3$  hold?

**Solution.** The trail  $y_1 - x_1 - x_2 - x_3 - x_4 - y_4$  is active: none of the nodes is in a collider configuration, so that their default state is open and conditioning on  $y_3$  does not block any of the nodes on the trail.

While  $x_1 - x_2 - x_3 - x_4$  forms a Markov chain, where e.g.  $x_4 \perp\!\!\!\perp x_1 \mid x_3$  holds, this not so for the distribution of the  $y$ 's.

### Exercise 8. *Alternative characterisation of independencies*

We have seen that  $x \perp\!\!\!\perp y|z$  is characterised by  $p(x, y|z) = p(x|z)p(y|z)$  or, equivalently, by  $p(x|y, z) = p(x|z)$ . Show that further equivalent characterisations are

$$p(x, y, z) = p(x|z)p(y|z)p(z) \quad \text{and} \quad (1)$$

$$p(x, y, z) = a(x, z)b(y, z) \quad \text{for some non-neg. functions } a(x, z) \text{ and } b(x, z). \quad (2)$$

The characterisation in Equation (2) will be important for undirected graphical models.

**Solution.** We first show the equivalence of  $p(x, y|z) = p(x|z)p(y|z)$  and  $p(x, y, z) = p(x|z)p(y|z)p(z)$ : By the product rule, we have

$$p(x, y, z) = p(x, y|z)p(z).$$

If  $p(x, y|z) = p(x|z)p(y|z)$ , it follows that  $p(x, y, z) = p(x|z)p(y|z)p(z)$ . To show the opposite direction assume that  $p(x, y, z) = p(x|z)p(y|z)p(z)$  holds. By comparison with the decomposition in the product rule, it follows that we must have  $p(x, y|z) = p(x|z)p(y|z)$  whenever  $p(z) > 0$  (it suffices to consider this case because for  $z$  where  $p(z) = 0$ ,  $p(x, y|z)$  may not be uniquely defined in the first place).

Equation (1) implies (2) with  $a(x, z) = p(x|z)$  and  $b(y, z) = p(y|z)p(z)$ . We now show the inverse. Let us assume that  $p(x, y, z) = a(x, z)b(y, z)$ . By the product rule, we have

$$p(x, y|z)p(z) = a(x, z)b(y, z). \quad (\text{S.50})$$

$$(\text{S.51})$$

Summing over  $y$  gives

$$\sum_y p(x, y|z)p(z) = p(z) \sum_y p(x, y|z) \quad (\text{S.52})$$

$$= p(z)p(x|z) \quad (\text{S.53})$$

Moreover

$$\sum_y p(x, y|z)p(z) = \sum_y a(x, z)b(y, z) \quad (\text{S.54})$$

$$= a(x, z) \sum_y b(y, z) \quad (\text{S.55})$$

so that

$$a(x, z) = \frac{p(z)p(x|z)}{\sum_y b(y, z)} \quad (\text{S.56})$$

Since the sum of  $p(x|z)$  over  $x$  equals one we have

$$\sum_x a(x, z) = \frac{p(z)}{\sum_y b(y, z)}. \quad (\text{S.57})$$

Now, summing  $p(x, y|z)p(z)$  over  $x$  yields

$$\sum_x p(x, y|z)p(z) = p(z) \sum_x p(x, y|z). \quad (\text{S.58})$$

$$= p(y|z)p(z) \quad (\text{S.59})$$

We also have

$$\sum_x p(x, y|z)p(z) = \sum_x a(x, z)b(y, z) \quad (\text{S.60})$$

$$= b(y, z) \sum_x a(x, z) \quad (\text{S.61})$$

$$\stackrel{(\text{S.57})}{=} b(y, z) \frac{p(z)}{\sum_y b(y, z)} \quad (\text{S.62})$$

so that

$$p(y|z)p(z) = p(z) \frac{b(y, z)}{\sum_y b(y, z)} \quad (\text{S.63})$$

We thus have

$$p(x, y, z) = a(x, z)b(y, z) \quad (\text{S.64})$$

$$\stackrel{(\text{S.56})}{=} \frac{p(z)p(x|z)}{\sum_y b(y, z)} b(y, z) \quad (\text{S.65})$$

$$= p(x|z)p(z) \frac{b(y, z)}{\sum_y b(y, z)} \quad (\text{S.66})$$

$$\stackrel{(\text{S.63})}{=} p(x|z)p(y|z)p(z) \quad (\text{S.67})$$

which is Equation (1).

### Exercise 9. More on independencies

*This exercise is on further properties and characterisations of statistical independence.*

- (a) *Without using  $d$ -separation, show that  $x \perp\!\!\!\perp \{y, w\} \mid z$  implies that  $x \perp\!\!\!\perp y \mid z$  and  $x \perp\!\!\!\perp w \mid z$ .*  
Hint: use the definition of statistical independence in terms of the factorisation of pmfs/pdfs.

**Solution.** We consider the joint distribution  $p(x, y, w|z)$ . By assumption

$$p(x, y, w|z) = p(x|z)p(y, w|z) \quad (\text{S.68})$$

We have to show that  $x \perp\!\!\!\perp y|z$  and  $x \perp\!\!\!\perp w|z$ . For simplicity, we assume that the variables are discrete valued. If not, replace the sum below with an integral.

To show that  $x \perp\!\!\!\perp y|z$ , we marginalise  $p(x, y, w|z)$  over  $w$  to obtain

$$p(x, y|z) = \sum_w p(x, y, w|z) \quad (\text{S.69})$$

$$= \sum_w p(x|z)p(y, w|z) \quad (\text{S.70})$$

$$= p(x|z) \sum_w p(y, w|z) \quad (\text{S.71})$$

Since  $\sum_w p(y, w|z)$  is the marginal  $p(y|z)$ , we have

$$p(x, y|z) = p(x|z)p(y|z), \quad (\text{S.72})$$


which means that  $x \perp\!\!\!\perp y|z$ .

To show that  $x \perp\!\!\!\perp w|z$ , we similarly marginalise  $p(x, y, w|z)$  over  $y$  to obtain  $p(x, w|z) = p(x|z)p(w|z)$ , which means that  $x \perp\!\!\!\perp w|z$ .

(b) For the directed graphical model below, show that the following two statements hold without using d-separation:

$$x \perp\!\!\!\perp y \quad \text{and} \tag{3}$$

$$x \not\perp\!\!\!\perp y \mid w \tag{4}$$



The exercise shows that not only conditioning on a collider node but also on one of its descendants activates the trail between  $x$  and  $y$ . You can use the result that  $x \perp\!\!\!\perp y \mid w \Leftrightarrow p(x, y, w) = a(x, w)b(y, w)$  for some non-negative functions  $a(x, w)$  and  $b(y, w)$ .

**Solution.** The graphical model corresponds to the factorisation

$$p(x, y, z, w) = p(x)p(y)p(z|x, y)p(w|z).$$

For the marginal  $p(x, y)$  we have to sum (integrate) over all  $(z, w)$

$$p(x, y) = \sum_{z, w} p(x, y, z, w) \tag{S.73}$$

$$= \sum_{z, w} p(x)p(y)p(z|x, y)p(w|z) \tag{S.74}$$

$$= p(x)p(y) \sum_{z, w} p(z|x, y)p(w|z) \tag{S.75}$$

$$= p(x)p(y) \underbrace{\sum_z p(z|x, y)}_1 \underbrace{\sum_w p(w|z)}_1 \tag{S.76}$$

$$= p(x)p(y) \tag{S.77}$$

Since  $p(x, y) = p(x)p(y)$  we have  $x \perp\!\!\!\perp y$ .

For  $x \not\perp\!\!\!\perp y \mid w$ , compute  $p(x, y, w)$  and use the result  $x \perp\!\!\!\perp y \mid w \Leftrightarrow p(x, y, w) = a(x, w)b(y, w)$ .

$$p(x, y, w) = \sum_z p(x, y, z, w) \tag{S.78}$$


$$= \sum_z p(x)p(y)p(z|x, y)p(w|z) \tag{S.79}$$

$$= p(x)p(y) \underbrace{\sum_z p(z|x, y)p(w|z)}_{k(x, y, w)} \tag{S.80}$$

Since  $p(x, y, w)$  cannot be factorised as  $a(x, w)b(y, w)$ , the relation  $x \perp\!\!\!\perp y \mid w$  cannot generally hold.

### Exercise 10. *Independencies in directed graphical models*

Consider the following directed acyclic graph.



For each of the statements below, determine whether it holds for all probabilistic models that factorise over the graph. Provide a justification for your answer.

(a)  $p(x_7|x_2) = p(x_7)$

**Solution.** Yes, it holds.  $x_2$  is a non-descendant of  $x_7$ ,  $\text{pa}(x_7) = \emptyset$ , and hence, by the local Markov property,  $x_7 \perp\!\!\!\perp x_2$ , so that  $p(x_7|x_2) = p(x_7)$ .

(b)  $x_1 \perp\!\!\!\perp x_3$

**Solution.** No, does not hold.  $x_1$  and  $x_3$  are d-connected, which only implies independence for *some* and not all distributions that factorise over the graph. The graph generally only allows us to read out independencies and not dependencies.

(c)  $p(x_1, x_2, x_4) \propto \phi_1(x_1, x_2)\phi_2(x_1, x_4)$  for some non-negative functions  $\phi_1$  and  $\phi_2$ .

**Solution.** Yes, it holds. The statement is equivalent to  $x_2 \perp\!\!\!\perp x_4 \mid x_1$ . There are three trails from  $x_2$  to  $x_4$ , which are all blocked:

1.  $x_2 - x_1 - x_4$ : this trail is blocked because  $x_1$  is in a tail-tail connection and it is observed, which closes the node.
2.  $x_2 - x_3 - x_6 - x_5 - x_4$ : this trail is blocked because  $x_3, x_6, x_5$  is in a collider configuration, and  $x_6$  is not observed (and it does not have any descendants).
3.  $x_2 - x_3 - x_6 - x_8 - x_7 - x_4$ : this trail is blocked because  $x_3, x_6, x_8$  is in a collider configuration, and  $x_6$  is not observed (and it does not have any descendants).

Hence, by the global Markov property (d-separation), the independency holds.

(d)  $x_2 \perp\!\!\!\perp x_9 \mid \{x_6, x_8\}$

**Solution.** No, does not hold. Conditioning on  $x_6$  opens the collider node  $x_4$  on the trail  $x_2 - x_1 - x_4 - x_7 - x_9$ , so that the trail is active.

(e)  $x_8 \perp\!\!\!\perp \{x_2, x_9\} \mid \{x_3, x_5, x_6, x_7\}$


**Solution.** Yes, it holds.  $\{x_3, x_5, x_6, x_7\}$  is the Markov blanket of  $x_8$ , so that  $x_8$  is independent of remaining nodes given the Markov blanket.

(f)  $\mathbb{E}[x_2 \cdot x_3 \cdot x_4 \cdot x_5 \cdot x_8 \mid x_7] = 0$  if  $\mathbb{E}[x_8 \mid x_7] = 0$

**Solution.** Yes, it holds.  $\{x_2, x_3, x_4, x_5\}$  are non-descendants of  $x_8$ , and  $x_7$  is the parent of  $x_8$ , so that  $x_8 \perp\!\!\!\perp \{x_2, x_3, x_4, x_5\} \mid x_7$ . This means that  $\mathbb{E}[x_2 \cdot x_3 \cdot x_4 \cdot x_5 \cdot x_8 \mid x_7] = \mathbb{E}[x_2 \cdot x_3 \cdot x_4 \cdot x_5 \mid x_7] \mathbb{E}[x_8 \mid x_7] = 0$ .

**Exercise 11. Independencies in directed graphical models**

Consider the following directed acyclic graph:



For each of the statements below, determine whether it holds for all probabilistic models that factorise over the graph. Provide a justification for your answer.

(a)  $x_1 \perp\!\!\!\perp x_2$

**Solution.** Does not hold. The trail  $x_1 - \theta_1 - \theta_2 - x_2$  is active (unblocked) because none of the nodes is in a collider configuration or in the conditioning set.

(b)  $p(x_1, y_1, \theta_1, u_1) \propto \phi_A(x_1, \theta_1, u_1) \phi_B(y_1, \theta_1, u_1)$  for some non-negative functions  $\phi_A$  and  $\phi_B$

**Solution.** Holds. The statement is equivalent to  $x_1 \perp\!\!\!\perp y_1 \mid \{\theta_1, u_1\}$ . The conditioning set  $\{\theta_1, u_1\}$  blocks all trails from  $x_1$  to  $y_1$  because they are both only in serial configurations in all trails from  $x_1$  to  $y_1$ , hence the independency holds by the global Markov property. Alternative justification: the conditioning set is the Markov blanket of  $x_1$ , and  $x_1$  and  $y_1$  are not neighbours which implies the independency.

(c)  $v_2 \perp\!\!\!\perp \{u_1, v_1, u_2, x_2\} \mid \{m_2, s_2, y_2, \theta_2\}$

**Solution.** Holds. The conditioning set is the Markov blanket of  $v_2$  (the set of parents, children, and co-parents): the set of parents is  $\text{pa}(v_2) = \{m_2, s_2\}$ ,  $y_2$  is the only child of  $v_2$ , and  $\theta_2$  is the only other parent of  $y_2$ . And  $v_2$  is independent of all other variables given its Markov blanket.

(d)  $\mathbb{E}[m_2 \mid m_1] = \mathbb{E}[m_2]$



**Solution.** Holds. There are four trails from  $m_1$  to  $m_2$ , namely via  $x_1$ , via  $y_1$ , via  $x_2$ , via  $y_2$ . In all trails the four variables are in a collider configuration, so that each of the trails is blocked. By the global Markov property (d-separation), this means that  $m_1 \perp\!\!\!\perp m_2$  which implies that  $\mathbb{E}[m_2 \mid m_1] = \mathbb{E}[m_2]$ .

Alternative justification 1:  $m_2$  is a non-descendent of  $m_1$  and  $\text{pa}(m_2) = \emptyset$ . By the directed local Markov property, a variable is independent from its non-descendants given the parents, hence  $m_2 \perp\!\!\!\perp m_1$ .

Alternative justification 2: We can choose a topological ordering where  $m_1$  and  $m_2$  are the first two variables. Moreover, their parent sets are both empty. By the directed ordered Markov, we thus have  $m_1 \perp\!\!\!\perp m_2$ .